

**TEKST NR 387**

**2000**

**A History of the Minimax Theorem:  
von Neumann's Conception  
of the Minimax Theorem -- a Journey  
Through Different Mathematical Contexts**

**Tinne Hoff Kjeldsen**

**TEKSTER fra**

**IMFUFA**

**ROSKILDE UNIVERSITETSCENTER**  
INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES  
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

IMFUFA, Roskilde University, Postbox 260, DK-4000 Roskilde, Denmark.

**A History of the Minimax Theorem: von Neumann's Conception of the Minimax Theorem – a Journey Through Different Mathematical Contexts**

by

**Tinne Hoff Kjeldsen**

IMFUFA text no. 387/2000

39 pages

ISSN 0106-6242

---

**Abstract:**

The purposes of this paper is first to tell the history of John von Neumann's development of the minimax theorem for two-person zero-sum games from his first proof of the theorem in 1928 until 1944 where he gave a completely different proof in the first coherent book on game theory. I will argue that von Neumann's conception of this theorem as a theorem belonging to the theory of linear inequalities as well as his awareness of its connection to fixpoint theorems were not present in 1928. In contradiction to the impression given in the literature these connections were only gradually recognized by von Neumann over time. By reading this knowledge into von Neumann's first proof of the minimax theorem from 1928 a major part of the cognitive development of this theorem is neglected within the history of mathematics. The significance of interactions between different branches of mathematics for the conception and development of the minimax theorem are neglected as well. My argumentation are based on an analysis of von Neumann's 1928-paper, of a paper he published in 1937 on a mathematical model for an expanding economy, and of the proof of the minimax theorem that appeared in von Neumann and Morgenstern's famous book on game theory published in 1944.

The second purpose of this paper is to discuss a more philosophical issue concerning the significance of the context in which a theorem is developed. The point of departure for this discussion is a dispute in 1953 between von Neumann and the French mathematician Maurice Fréchet about who should be named the initiator of game theory – an honour the mathematical literature at that time associated with von Neumann. Fréchet argued that eventhough Émile Borel was not able to prove the minimax theorem he was the true initiator of game theory due to his papers on the subject published in the beginning of the twenties before von Neumann's 1928 paper. The interesting issue here is not to settle the priority between Borel and von Neumann but rather to analyze the significance of the minimax theorem.

# A History of the Minimax Theorem: von Neumann's Conception of the Minimax Theorem – a Journey Through Different Mathematical Contexts

Tinne Hoff Kjeldsen, Department of Mathematics, Roskilde University, P.O. Box 260, DK-4000 Roskilde, Denmark.

## 1 Introduction

The purposes of this paper is first to tell the history of John von Neumann's development of the minimax theorem for two-person zero-sum games from his first proof of the theorem in 1928 until 1944 where he gave a completely different proof in the first coherent book on game theory. I will argue that von Neumann's conception of this theorem as a theorem belonging to the theory of linear inequalities as well as his awareness of its connection to fixpoint theorems were not present in 1928. In contradiction to the impression given in the literature these connections were only gradually recognized by von Neumann over time. By reading this knowledge into von Neumann's first proof of the minimax theorem from 1928 a major part of the cognitive development of this theorem is neglected within the history of mathematics. The significance of interactions between different branches of mathematics for the conception and development of the minimax theorem are neglected as well. This paper will remedy this and shed new light on these issues.

Since the beginning of the nineties there has been an increasing interest in the history of game theory, several historical papers have appeared and most of them of course mention von Neumann's 1928 proof of the minimax theorem. A common feature though is that none of these give an analysis of

the mathematics in von Neumann's proof. There is only one paper that goes deeper into the mathematics. It is an older essay written by two Princeton mathematicians; the late Albert W. Tucker and Harold W. Kuhn in memory of John von Neumann. They treat the mathematics in a modern (-1958) framework and emphasize in particular the connections to fixpoint theorems and the theory of linear inequalities [Kuhn and Tucker, 1958, p. 111-112]. The other historical papers state about von Neumann's 1928 proof of the minimax theorem that it is very difficult.<sup>1</sup> The von Neumann biographer Steve J. Heims very telling called it "a tour de force" [Heims, 1980, p. 91]. Some of the papers also state that the proof is about systems of linear inequalities and equations<sup>2</sup> and one that it is based on fixpoint theorems<sup>3</sup>. Reading von Neumann's 1928 paper I found that these statements were not at all obvious, as a matter of fact von Neumann did not talk about fixpoints in his 1928 proof and he did not formulate or present a system of linear inequalities and equations to be solved. Today we know that all these connections are there but I think that von Neumann was not fully aware of that in 1928, it was an insight that emerged gradually during his work from 1928 until 1944 in which the minimax theorem – some times surprisingly – presented itself in different mathematical contexts.

I will argue for this claim through an analysis of von Neumann's 1928-paper, of a paper he published in 1937 on a mathematical model for an expanding economy, and of the proof of the minimax theorem that appeared in von Neumann and Morgenstern's famous book on game theory published in 1944.

The second purpose of this paper is to discuss a more philosophical issue concerning the significance of the context in which a theorem is developed. The importance of a mathematical theorem is dependent on the branch or discipline of mathematics within which it is considered. A mathematical result is not likely to be deemed equally important within different branches or contexts of mathematics. The interesting questions, the questions that guide the research in different mathematical contexts are not the same. Thus, the potential of a mathematical theorem for stimulating further research is dependent of the mathematical context of discovery.<sup>4</sup>

---

<sup>1</sup>See [Dimand and Dimans, 1992, p. 24], [Leonard, 1992, p. 44], [Ingrao and Israel, 1990, p. 211].

<sup>2</sup>See [Ingrao and Israel, 1990, p. 211], [Heims, 1980, p. 91].

<sup>3</sup>See [Ingrao and Israel, 1990, p. 211].

<sup>4</sup>For example the calculus of variations with constraints and mathematical program-

The background for these questions in relation to the history of the minimax theorem is a dispute in 1953 between von Neumann and the French mathematician Maurice Fréchet about who should be named the initiator of game theory – an honour the mathematical literature at that time associated with von Neumann. Fréchet argued that eventhough Émile Borel was not able to prove the minimax theorem he was the true initiator of game theory due to his papers on the subject published in the beginning of the twenties before von Neumann's 1928 paper. The interesting issue is not to settle the priority between Borel and von Neumann but rather to analyze the significance of the minimax theorem. According to Fréchet the minimax theorem was not such an important result because it turned out that it can be derived very easily from other theorems on linear inequalities, theorems proved before 1928. The underlying assumption behind Fréchet's argumentation is that theorems that turn out to be equivalent have the same significance or the same potential for stimulating further mathematical developments regardless of the mathematical context in which they were derived. This touches a very interesting philosophical issue I think, namely the significance of the mathematical context for which kind of new questions of investigation a theorem can give rise to. A contextualized analysis of similar mathematical theorems derived in different mathematical contexts can give the historian a tool for understanding mechanisms behind the division of mathematical results that gave rise to new developments in mathematics and results that did not.<sup>5</sup>

ming both treat optimization under constraints. But in the calculus of variation the infinite cases are treated where as mathematical programming is concerned with finite dimensional cases, so a theorem about constrained optimization can be deemed very important and can lead to new knowledge in mathematical programming where as the same theorem evaluated from the point of view of the calculus of variation is seen as just a minor thing. To be specific this was the case for the so-called Kuhn-Tucker theorem in mathematical programming, see [Kjeldsen, 2000].

<sup>5</sup>The Kuhn-Tucker theorem in nonlinear programming is an example of this. Kuhn and Tucker derived the theorem in 1950 and it immediately launched the theory of nonlinear programming and is viewed as a very important result. Later it turned out that a similar result had been proven only 11 years earlier by William Karush in his master thesis. Karush's work was done in the mathematical context of the calculus of variations within which it was not regarded as a very important result, it was not even published. Also Fritz John proved a similar result which he had problems getting published. It finally appeared on print in 1948 – only two years before Kuhn and Tucker's version of the theorem was published. John's work was done within the context of the theory of convexity in which the theorem was not deemed to be something special. The reasons for the very different receptions of these results within the mathematical community can be explained by referring

This issue is discussed in the second part of the paper on the basis of an analysis of some of Borel's papers on game theory, of von Neumann's work, and of the dispute between Fréchet and von Neumann.

## 2 The First Proof of the Minimax Theorem: von Neumann's 1928 Paper

John (Johann) von Neumann (1903-1957) published his first paper on what he called "Theorie der Gesellschaftsspiele" in 1928. From there on it took 16 years before von Neumann published on game theory again, so for a long time the 1928-paper stood as a kind of singularity in his mathematical production. This has of course given rise to some speculations about why and where this idea and inspiration to develop a mathematical theory of games came from. Two explanations are suggested in the literature, one of them is that von Neumann got the idea from reading Borel's work on the subject.<sup>6</sup> Von Neumann himself claimed that he developed the theory independently of Borel [von Neumann, 1928a]. In the 1928-paper von Neumann has a footnote telling that someone drew his attention to the notes of Borel during the proofreading [von Neumann, 1928, p. 306]. I think von Neumann's claim is supported by the course of events: Already in December 1926 he presented his work at the weekly seminar of the mathematical institute in Göttingen [von Neumann, 1928, p. 295]. Eventhough he sent the manuscript to *Mathematische Annalen* in July 1927 it was not until May 1928, that is allmost a year later, that he had Borel present the work including the minimax theorem to the Académie des Sciences in Paris, just a short time before the paper itself was published [von Neumann, 1928a]. When he heard about Borel's work he might have been afraid that Borel or someone else would be about to publish a similar result and then acted quickly at that time by sending a note to the Académie to ensure priority. The other explanation for why von Neumann suddently developed a theory of games is more plausible I think. Here the apperance of von Neumann's game theoretic work is linked with the social context of von Neumann's life during the years leading up to the publication of the paper. Von Neumann was at that time very

---

to the significance of the different mathematical contexts in which the results were derived. For an analysis of this see [Kjeldsen, 2000].

<sup>6</sup>See for example [Ulam, 1958, p. 7].

much influenced by Hilbert and the Göttingen mathematical community. He was especially deeply involved in Hilbert's axiomatization programme.<sup>7</sup> In the papers [Mirowski, 1991, 1992] Mirowski argues convincingly that von Neumann's game theory was a result of his connection to Hilbert and the formalist programme.<sup>8</sup>

## 2.1 What is a "Gesellschaftsspiele"?

The two essential parts of von Neumann's 1928 paper are the mathematization of "Gesellschaftsspiele" or "games of strategy" and the proof of the theorem "Max Min=Min Max" for a game involving two players who play against each other and for which the players total gain add up to zero. That is the theorem now known as the minimax theorem for two-person zero-sum games. In the following I will explain how von Neumann mathematized games of strategy and how he proved the minimax theorem.<sup>9</sup>

Von Neumann began the paper by posing the question under consideration

*n* Spieler,  $S_1, S_2, \dots, S_n$ , spielen ein gegebenes Gesellschaftsspiel  $B$ . Wie muß einer dieser Spieler,  $S_m$ , spielen, um dabei ein möglichst günstiges Resultat zu erzielen?<sup>10</sup> [von Neumann, 1928, p. 295]

As von Neumann pointed out the problem is well known from daily life but ambiguous because what will happen when there are more than player involved? In that case the fate of each player depends on the rest of the players and they are all guided by the same selfish interests. Thus the first problem von Neumann faced was to clarify what precisely was to be understood by the term "Gesellschaftsspiel". As the following quote shows von Neumann had a very broad understanding of the concept

<sup>7</sup>In 1925 - 1928 he published three papers on the axiomatization of set theory, one on Hilbert's proof theory, and seven papers on the foundation and axiomatization of quantum mechanics; see the bibliography of John von Neumann in his collected works [von Neumann, 1963, pp. 645 - 652].

<sup>8</sup>See also [Leonard, 1992, 1995].

<sup>9</sup>The paper was published in German. In 1959 an English translation of it was published from which the translations in the footnotes of the quotes have been taken.

<sup>10</sup>" $n$  players  $S_1, S_2, \dots, S_n$  are playing a game of strategy,  $B$ . How must one of the participants,  $S_m$ , play in order to achieve a most advantageous result?" [von Neumann, 1928, (1959 p. 13)].

Es fallen unter diesen Begriff sehr viele, recht verschiedenartige Dinge: von der Roulette bis zum Schach, vom Bakkarat bis zum Bridge liegen ganz verschiedene Varianten des Sammelbegriffes 'Gesellschaftsspiel' vor. Und letzten Endes kann auch irgendein Ereignis, mit gegebenen äusseren Bedingungen und gegebenen Handelnden (den absolut freien Willen der letzteren vorausgesetzt), als Gesellschaftsspiel angesehen werden, wenn man seine Rückwirkungen auf die in ihm handelnden Personen betrachtet.<sup>11</sup> [von Neumann, 1928, p. 295]

One must say that von Neumann's very broad interpretation of "Gesellschaftsspiele" points towards an extremely ambitious project. At a first glance it must have seemed very unlikely that one could succeed in building a mathematical model for this kind of situation. Anyway even though von Neumann was 'only' able to construct a solution concept and prove the existence of such a solution for a very limited subset of the overall game concept, he started out with the mathematization of the general case.

By collecting the common features in game situations von Neumann derived a qualitative description of the game concept. He argued as follows: A game is composed of a series of events of which each can have at most a finite number of outcomes. In some game situations it can happen that the outcome of some of the events depends only on chance. This means that the probabilities with which each of the outcomes will appear are known but none of the players have any influence on them. The outcome of all other events are subject to the individual player's free choices. For each of these events it is known which player determines the outcome, and what kind of information this player has regarding the outcome of earlier events. Finally there is a rule by which the winnings and losses of each player can be calculated after the game, that is after the outcome of all events in the play are known. [von Neumann, 1928, p. 296].

In order to be able to work with this very broad concept of a game von Neumann reformulated the above qualitative description in a more precise form which then served as his definition of a game of strategy. His definition

---

<sup>11</sup> "A great many different things come under this heading, anything from roulette to chess, from baccarat to bridge. And after all, any event – given the external conditions and the participants in the situation (provided the latter are acting of their own free will) – may be regarded as a game of strategy if one looks at the effect it has on the participants." [von Neumann, 1928 (1959 p. 13)].

was build up around five points.

The first one specifies the number ( $z$ ) of events depending on chance and the number ( $s$ ) of events depending on the free will of the players. Von Neumann let

$$E_1, E_2, \dots, E_z$$

denote the events depending on chance, and

$$F_1, F_2, \dots, F_s$$

denote the events depending on the free will of the players.

The second is the specification of the number  $M_\mu$  ( $\mu = 1, 2, \dots, z$ ) of possible outcomes of each single event of chance  $E_\mu$ , and the number  $N_\nu$  ( $\nu = 1, 2, \dots, s$ ) of possible outcomes of each single event of free will  $F_\nu$ . Von Neumann referred to a result of an event by its number, i. e.  $1, 2, \dots, M_\mu$  or  $1, 2, \dots, N_\nu$ .

The third thing one needs to know in von Neumann's game model is the probabilities  $\alpha_\mu^{(1)}, \alpha_\mu^{(2)}, \dots, \alpha_\mu^{(M_\mu)}$  with which the outcomes  $1, 2, \dots, M_\mu$  of an event of chance  $E_\mu$  will occur, thus

$$\alpha_\mu^{(1)} \geq 0, \alpha_\mu^{(2)} \geq 0, \dots, \alpha_\mu^{(M_\mu)} \geq 0,$$

and

$$\alpha_\mu^{(1)} + \alpha_\mu^{(2)} + \dots + \alpha_\mu^{(M_\mu)} = 1.$$

For every event of free will  $F_\nu$ , one also needs to specify which player  $S_m$  determines the outcome of this event and in addition one also needs to know what have occurred up to this moment, that is the corresponding numbers for all earlier events both those of chance and those of free will that the player in charge have information about when he or she make up his or her mind.

Finally one needs to specify  $n$  real valued functions  $f_1, f_2, \dots, f_n$  of  $z + s$  variables, where the first  $z$  ariables can take the values

$$1, 2, \dots, M_1; \quad 1, 2, \dots, M_2; \quad \dots; \quad 1, 2, \dots, M_z;$$

and the last  $s$  variables can take the values

$$1, 2, \dots, N_1; \quad 1, 2, \dots, N_2; \quad \dots; \quad 1, 2, \dots, N_s.$$

These functions determine the gain of the players and must add up to zero

$$f_1 + f_2 + \dots + f_n \equiv 0.$$

Suppose the results of the  $z$  events of chance and the  $s$  events of free will in a game turned out to be

$$x_1, x_2, \dots, x_z, \quad y_1, y_2, \dots, y_s,$$

respectively, where

$$x_\mu \in \{1, 2, \dots, M_\mu\}, \quad y_\nu \in \{1, 2, \dots, N_\nu\},$$

$$\mu = 1, 2, \dots, z, \quad \nu = 1, 2, \dots, s,$$

the players  $S_1, S_2, \dots, S_n$  then 'gain' the amounts

$$f_1(x_1, \dots, x_z, y_1, \dots, y_s), f_2(x_1, \dots, x_z, y_1, \dots, y_s), \dots, \\ f_n(x_1, \dots, x_z, y_1, \dots, y_s)$$

from each other [von Neumann, 1928, p. 296 - 297].

The above definition was von Neumann's definition of a game of strategy. But as he himself pointed out the notion of a player  $S_m$ , trying to achieve a result as advantageous as possible is kind of obscure. It is clear that the most advantageous result for  $S_m$  has to be defined as the largest possible value of  $f_m$ , but  $f_m$  depends of  $z + s$  variables of which only a part is controlled by  $S_m$ , and this is exactly the heart of the problem:

Es soll versucht werden, die Rückwirkungen der Spieler aufeinander zu untersuchen, die Konsequenzen des (für alles soziale Geschehen so charakteristischen!) Umstandes, daß jeder Spieler auf die Resultate aller anderen einen Einfluß hat und dabei nur am eigenen interessiert ist.<sup>12</sup> [von Neumann, 1928, p. 298]

The next step in von Neumann's building of the theory was to simplify the game concept as much as possible without losing anything in generality. The key trick was the introducing of the concept of strategy by which he could reduce the number of events of free will to the number of players, such

<sup>12</sup>"We shall try to investigate the effects which the players have on each other, the consequences of the fact (so typical of all social happenings!) that each player influences the results of all other players, even though he is only interested in his own [von Neumann, 1928, (1959 p. 17)].

that event number  $\nu$  is determined by the free will of player  $S_\nu$ . All the information about the other players decisions and the outcome of the events of chance that a player has access to is inherent in the concept of strategy. The consequence of this is, that each player choose his or her strategy without knowing neither the strategies chosen by the others nor the results of the events of chance.

Another advantage of the concept of strategy is, that von Neumann could eliminate the events of chance all together. First he reduced the number of events of chance to one. Because, since a player has to choose the strategy without knowing beforehand the outcome of the events of chance, these events need no longer be treated as separate events. It is then possible to combine all  $z$  events of chance into one single event of chance  $H$ , the outcome of which will be a collection of numbers

$$x_1, x_2, \dots, x_z \quad (x_\mu = 1, 2, \dots, M_\mu, \mu = 1, 2, \dots, z),$$

with their respective probabilities

$$\alpha_1^{(x_1)} \alpha_2^{(x_2)} \dots \alpha_z^{(x_z)},$$

There are  $M = M_1 M_2 \dots M_z$  possible collections of these numbers. Von Neumann associated each collection with a number

$$1, 2, \dots, M \quad (M = M_1 M_2 \dots M_z),$$

and he let

$$\beta_1, \beta_2, \dots, \beta_M$$

denote the corresponding probabilities [von Neumann, 1928, p. 300].

By doing so von Neumann had it all boiled down to the following: If the players  $S_1, S_2, \dots, S_n$  have chosen the strategies

$$S_{u_1}^{(1)}, S_{u_2}^{(2)}, \dots, S_{u_n}^{(n)},$$

where

$$u_m = 1, 2, \dots, \Sigma_m, \quad m = 1, 2, \dots, n,$$

and if the outcome of the event of chance  $H$ , is the number  $\nu (= 1, 2, \dots, M)$ , then the results for the players  $S_1, S_2, \dots, S_n$  are

$$f_1(\nu, u_1, u_2, \dots, u_n), f_2(\nu, u_1, u_2, \dots, u_n), \dots, f_n(\nu, u_1, u_2, \dots, u_n)$$

respectively.<sup>13</sup> Now if only the choices  $u_1, u_2, \dots, u_n$ , and not the result  $\nu$ , of the event of chance is known, then the expected value of  $f_1, f_2, \dots, f_n$  would be

$$g_m(u_1, \dots, u_n) = \sum \beta_\nu f_m(\nu, u_1, \dots, u_n), \quad (m = 1, 2, \dots, n),$$

( $f_1 + \dots + f_n \equiv 0$  implies  $g_1 + \dots + g_n \equiv 0$ ). Von Neumann then argued that according to the theory of probability it is fully acceptable to ignore the event of chance and instead work with the expected values  $g_1, \dots, g_n$ . That is, by substituting the *exact* results ( $f_m$ ) for the individual players by the *expected* values he eliminated  $H$  all together.

These simplifications left von Neumann with the following formulation of a game of strategy: Each of the players  $S_1, S_2, \dots, S_n$  choose a number without any information about the choice of the others.  $S_m$  chooses among the numbers  $1, 2, \dots, \Sigma_m$ . After the choices  $x_1, x_2, \dots, x_n$  ( $x_m = 1, 2, \dots, \Sigma_m, m = 1, 2, \dots, n$ ) have been made the players will receive the amount

$$g_1(x_1, \dots, x_n), g_2(x_1, \dots, x_n), \dots, g_n(x_1, \dots, x_n),$$

respectively, where  $g_1 + \dots + g_n = 0$  holds [von Neumann, 1928, p. 301 - 302].

## 2.2 The case n=2

Von Neumann was in 1928 not able to prove anything according to the existence of optimal strategies for the general case. In order to do so he analyzed the simplest case, namely a game of strategy with only two players  $S_1$  and  $S_2$ . The situation is then, that the player  $S_1$  choose a number  $x \in \{1, 2, \dots, \Sigma_1\}$ ,  $S_2$  choose a number  $y \in \{1, 2, \dots, \Sigma_2\}$  without knowing what the other player has chosen, and they then receive the amount  $g(x, y), -g(x, y)$  respectively. Von Neumann then posed the question under consideration:

Es ist leicht, sich ein Bild von den Tendenzen zu machen, die in einem solchen 2-Personen-Spiele miteinander kämpfen: Es wird von zwei Seiten am Werte von  $g(x, y)$  hin und her gezerzt, nämlich durch  $S_1$ , der ihn möglichst gross, und durch  $S_2$ , der ihn möglichst klein machen will.  $S_1$ , gebietet über die Variable  $x$ , und  $S_2$  über

<sup>13</sup>Earlier von Neumann had argued that each player  $S_m$  only have a finite number of strategies ( $S_1^m, S_2^m, \dots, S_{\sigma_m}^m$ ) to choose from.

die Variable  $y$ . Was wird geschehen?<sup>14</sup> [von Neumann, 1928, p. 302]

The core question is 'What will happen?' Von Neumann's analysis of the situation ran as follows: If  $S_1$  chose the number  $x_0$  ( $x_0 = 1, 2, \dots, \Sigma_1$ ), that is the strategy  $x_0$ , his result  $g(x_0, y)$  would then also depend on the choice of  $S_2$  but no matter which choice ( $y$ )  $S_2$  comes up with the following inequality will be true

$$g(x_0, y) \geq \min_y g(x_0, y).$$

Now if we suppose (against the rules of the game) that  $S_2$  knew  $x_0$ ,  $S_2$  would according to the assumptions in the model choose  $y = y_0$  such that

$$g(x_0, y_0) = \min_y g(x_0, y).$$

Facing this situation the best thing for  $S_1$  would be to choose  $x_0$  such that

$$\min_y g(x_0, y) = \max_x \min_y g(x, y).$$

The conclusion of von Neumann is then that  $S_1$  can make

$$g(x_0, y) \geq \max_x \min_y g(x, y),$$

independently of the choice of  $S_2$ . The same argument holds for  $S_2$ , which can make

$$g(x, y_0) \leq \min_y \max_x g(x, y),$$

no matter what strategy  $x$ ,  $S_1$  chooses.

From this von Neumann concluded that if a pair of strategies  $x_0, y_0$  can be found for which

$$g(x_0, y_0) = \max_x \min_y g(x, y) = \min_y \max_x g(x, y) = M,$$

then that would necessary be the choices for  $S_1$  and  $S_2$  respectively, and  $M$  would be the result of the game [von Neumann, 1928, pp. 302 - 303]. Thus, such a pair of strategies  $x_0, y_0$  if they exist would constitute a solution

<sup>14</sup> "It is easy to picture the forces struggling with each other in such a two-person game. The value of  $g(x, y)$  is being tugged at from two sides, by  $S_1$  who wants to maximize it, and by  $S_2$  who wants to minimize it.  $S_1$  controls the variable  $x$ ,  $S_2$  the variable  $y$ . What will happen?" [von Neumann, 1928, (1959 p. 21)].

concept for two-person games. Unfortunately the existence of such a pair of strategies is not automatically guaranteed.

The trick used by von Neumann to overcome this difficulty was to introduce what is now known as mixed strategies. Instead of choosing an  $x$  or a  $y$ , the players specify the probabilities with which they will choose the different strategies. That is, the player  $S_1$  chooses  $\Sigma_1$  probabilities

$$\xi_1, \xi_2, \dots, \xi_{\Sigma_1} \quad (\xi_1 \geq 0, \xi_2 \geq 0, \dots, \xi_{\Sigma_1} \geq 0, \quad \sum \xi_i = 1),$$

and from an urn containing the numbers  $1, 2, \dots, \Sigma_1$  with the above specified probabilities, he or she draws a number and chooses that number. Analog  $S_2$  specifies  $\Sigma_2$  probabilities

$$\eta_1, \eta_2, \dots, \eta_{\Sigma_2} \quad (\eta_1 \geq 0, \eta_2 \geq 0, \dots, \eta_{\Sigma_2} \geq 0, \quad \sum \eta_j = 1).$$

Von Neumann put

$$(\xi_1, \xi_2, \dots, \xi_{\Sigma_1}) = \xi, \text{ and } (\eta_1, \eta_2, \dots, \eta_{\Sigma_2}) = \eta.$$

If  $S_1$  chooses  $\xi$ , and  $S_2$  chooses  $\eta$ , the expected value of the amount  $S_1$  receives is

$$h(\xi, \eta) = \sum_{p=1}^{\Sigma_1} \sum_{q=1}^{\Sigma_2} g(p, q) \xi_p \eta_q,$$

while the expected value for  $S_2$  is  $-h(\xi, \eta)$  [von Neumann, 1928, p. 304].

As before von Neumann argued that  $S_1$  is in a position to obtain the minimal expected value  $\max_{\xi} \min_{\eta} h(\xi, \eta)$  no matter what  $S_2$  chooses to do.  $S_2$  can keep the expected value of  $S_1$  from exceeding the maximal value  $\min_{\eta} \max_{\xi} h(\xi, \eta)$ . By considering the mixed strategies instead of pure strategies the expected values of the players is expressed by the bilinear form  $h$ , and for those von Neumann was able to show that there always exist mixed strategies  $\xi_0, \eta_0$  such that

$$\max_{\xi} \min_{\eta} h(\xi, \eta) = \min_{\eta} \max_{\xi} h(\xi, \eta) = h(\xi_0, \eta_0).$$

This result is the famous minimax theorem of von Neumann and it establishes that for two-person games of this kind there always exist optimal (mixed-) strategies. This is called the minimax solution concept of two-person zero-sum games. It has been criticized for being too defensive a solution concept, indeed it is a solution telling you what is the best you can do in the worst possible case.

## 2.3 Von Neumann's proof of the minimax theorem

Actually von Neumann proved a generalized version of the minimax theorem. He considered a broader class of functions than the bilinear forms  $h$ , and formulated the theorem in the following way:

For continuous functions  $f$  of two variables  $\xi \in \mathbf{R}^M$ ,  $\eta \in \mathbf{R}^N$ ,  $\xi \geq 0$ ,  $\eta \geq 0$ ,  $\xi_1 + \dots + \xi_M \leq 1$ ,  $\eta_1 + \dots + \eta_N \leq 1$  satisfying the condition:

(K.) Wenn  $f(\xi', \eta) \geq A$ ,  $f(\xi'', \eta) \geq A$  ist, so ist auch für jedes  $0 \leq \nu \leq 1$ ,  $\xi = \nu\xi' + (1 - \nu)\xi''$  (d.h.  $\xi_p = \nu\xi'_p + (1 - \nu)\xi''_p$ ,  $p = 1, 2, \dots, M$ )  $f(\xi, \eta) \geq A$ . Wenn  $f(\xi, \eta') \leq A$ ,  $f(\xi, \eta'') \leq A$  ist, so ist auch für jedes  $0 \leq \nu \leq 1$ ,  $\eta = \nu\eta' + (1 - \nu)\eta''$  (d.h.  $\eta_q = \nu\eta'_q + (1 - \nu)\eta''_q$ ,  $q = 1, 2, \dots, N$ )  $f(\xi, \eta) \leq A$ .

... Für diese Funktionen  $f(\xi, \eta)$  werden wir beweisen:

$$\max_{\xi} \min_{\eta} f(\xi, \eta) = \min_{\eta} \max_{\xi} f(\xi, \eta),$$

wobei  $\max_{\xi}$  über  $\xi_1 \geq 0, \dots, \xi_M \geq 0$ ,  $\xi_1 + \dots + \xi_M \leq 1$ , und  $\min_{\eta}$  über  $\eta_1 \geq 0, \dots, \eta_N \geq 0$ ,  $\eta_1 + \dots + \eta_N \leq 1$  zu erstrecken ist.<sup>15</sup>  
[von Neumann, 1928, p. 307]

Today a function with the property (K) is called quasiconcave in  $\xi$  and quasiconvex in  $\eta$ .

Since the function  $h(\xi, \eta)$ , that determines the expected values for the two players is bilinear, it is also continuous and has the property (K), so a proof of this theorem will also prove the existence of optimal strategies for a two-person zero-sum game.

<sup>15</sup>“(K.) If  $f(\xi', \eta) \geq A$ ,  $f(\xi'', \eta) \geq A$ , then  $f(\xi, \eta) \geq A$  for every  $0 \leq \nu \leq 1$ ,  $\xi = \nu\xi' + (1 - \nu)\xi''$  (i.e.,  $\xi_p = \nu\xi'_p + (1 - \nu)\xi''_p$ ,  $p = 1, 2, \dots, M$ ). If  $f(\xi, \eta') \leq A$ ,  $f(\xi, \eta'') \leq A$ , then  $f(\xi, \eta) \leq A$  for every  $0 \leq \nu \leq 1$ ,  $\eta = \nu\eta' + (1 - \nu)\eta''$  (i.e.,  $\eta_q = \nu\eta'_q + (1 - \nu)\eta''_q$ ,  $q = 1, 2, \dots, N$ ).

... For these functions  $f(\xi, \eta)$  we are going to prove that

$$\max_{\xi} \min_{\eta} f(\xi, \eta) = \min_{\eta} \max_{\xi} f(\xi, \eta),$$

where  $\max_{\xi}$  is taken over the range  $\xi_1 \geq 0, \dots, \xi_M \geq 0$ ,  $\xi_1 + \dots + \xi_M \leq 1$  and  $\min_{\eta}$  is taken over the range  $\eta_1 \geq 0, \dots, \eta_N \geq 0$ ,  $\eta_1 + \dots + \eta_N \leq 1$ .” [von Neumann, 1928, (1959 p. 26-27)].

Von Neumann began by rewriting

$$\max_{\xi} \min_{\eta} f(\xi, \eta) = \min_{\eta} \max_{\xi} f(\xi, \eta)$$

in the form

$$\begin{aligned} & \max_{\substack{\xi_1 \geq 0 \\ \xi_1 \leq 1}} \max_{\substack{\xi_2 \geq 0 \\ \xi_1 + \xi_2 \leq 1}} \dots \max_{\substack{\xi_M \geq 0 \\ \xi_1 + \dots + \xi_M \leq 1}} \min_{\substack{\eta_1 \geq 0 \\ \eta_1 \leq 1}} \min_{\substack{\eta_2 \geq 0 \\ \eta_1 + \eta_2 \leq 1}} \dots \min_{\substack{\eta_N \geq 0 \\ \eta_1 + \dots + \eta_N \leq 1}} f(\xi, \eta) \\ & = \min_{\substack{\eta_1 \geq 0 \\ \eta_1 \leq 1}} \min_{\substack{\eta_2 \geq 0 \\ \eta_1 + \eta_2 \leq 1}} \dots \min_{\substack{\eta_N \geq 0 \\ \eta_1 + \dots + \eta_N \leq 1}} \max_{\substack{\xi_1 \geq 0 \\ \xi_1 \leq 1}} \max_{\substack{\xi_2 \geq 0 \\ \xi_1 + \xi_2 \leq 1}} \dots \max_{\substack{\xi_M \geq 0 \\ \xi_1 + \dots + \xi_M \leq 1}} f(\xi, \eta). \end{aligned}$$

By putting

$$M^{\xi_r} f(\xi_1, \dots, \xi_r, \eta_1, \dots, \eta_s) = \max_{\xi_r} f(\xi_1, \dots, \xi_r, \eta_1, \dots, \eta_s),$$

$$M^{\eta_s} f(\xi_1, \dots, \xi_r, \eta_1, \dots, \eta_s) = \min_{\eta_s} f(\xi_1, \dots, \xi_r, \eta_1, \dots, \eta_s),$$

he eliminated  $f$ 's dependency of  $\xi_r$  and  $\eta_s$  respectively. Thus von Neumann wrote the identity under consideration as

$$M^{\xi_1} M^{\xi_2} \dots M^{\xi_p} M^{\eta_1} M^{\eta_2} \dots M^{\eta_q} f = M^{\eta_1} M^{\eta_2} \dots M^{\eta_q} M^{\xi_1} M^{\xi_2} \dots M^{\xi_p} f.$$

With  $p = M$  and  $q = N$  this is equivalent with von Neumann's formulation above of the minimax theorem, where he considered  $\xi \in \mathbf{R}^M$  and  $\eta \in \mathbf{R}^N$ .

With these reformulations as a tool von Neumann reduced the proof to the proof of the following two lemmata:

$\alpha$ ) If  $f = f(\xi_1, \dots, \xi_r, \eta_1, \dots, \eta_s)$  is continuous and has the property **(K)** then  $M^{\xi_r} f$  and  $M^{\eta_s} f$  are continuous and fullfil **(K)**.

$\beta$ ) If  $f = f(\xi_1, \dots, \xi_r, \eta_1, \dots, \eta_s)$  is continuous and fullfil **(K)** then

$$M^{\xi_r} M^{\eta_s} f = M^{\eta_s} M^{\xi_r} f.$$

It is easy to see that the minimax theorem can be derived directly from  $\alpha$ ) and  $\beta$ ).

Von Neumann's proof of  $\alpha$ ) is straight forward. The continuity of  $M^{\xi_r} f$  and  $M^{\eta_s} f$  is a direct consequence of the continuity of  $f$ . To prove that  $M^{\xi_r} f$  and  $M^{\eta_s} f$  has the property **(K)** von Neumann used first that

a continuous function on a closed and boundet set has a maximum and a minimum and second that  $f$  itself has the property (K).<sup>16</sup>

The central part of the proof is  $\beta$ ) which is much more complicated to prove. In what follows I will go through the proof step by step following von Neumann closely. Then I will comment on the proof and discuss it in relation to the description given by Kuhn and Tucker.

### The proof for $\beta$ )

Von Neumann is going to prove that  $M^{\xi_r} M^{\eta_s} f = M^{\eta_s} M^{\xi_r} f$  for all  $\xi_1, \dots, \xi_{r-1}, \eta_1, \dots, \eta_{s-1}$ . He began by considering  $f$  for some fixed  $\xi_1, \dots, \xi_{r-1}, \eta_1, \dots, \eta_{s-1}$ . Then  $f$  is a function of  $\xi_r$  og  $\eta_s$  alone and  $f$  is obviously still continuous and posses the property (K). Writting  $\xi$  and  $\eta$  instead of  $\xi_r$  and  $\eta_s$  respectively what von Neumann is going to show is that

$$\max_{0 \leq \xi \leq a} \min_{0 \leq \eta \leq b} f(\xi, \eta) = \min_{0 \leq \eta \leq b} \max_{0 \leq \xi \leq a} f(\xi, \eta),$$

where  $a = 1 - \xi_1 - \dots - \xi_{r-1}$  and  $b = 1 - \eta_1 - \dots - \eta_{s-1}$ . As pointed out by von Neumann this can also be formulated in an other way:

Es gibt einen "Sattelpunkt"  $\xi_0, \eta_0$  ( $0 \leq \xi_0 \leq a, 0 \leq \eta_0 \leq b$ ), d.h.  $f(\xi_0, \eta)$  nimmt in  $0 \leq \eta \leq b$  sein Minimum für  $\eta = \eta_0$  an, und  $f(\xi, \eta_0)$  nimmt in  $0 \leq \xi \leq a$  sein Maximum für  $\xi = \xi_0$  an.<sup>17</sup> [von Neumann, 1928, p. 309].

Because it is always true that

$$\max_{\xi} \min_{\eta} f(\xi, \eta) \leq \min_{\eta} \max_{\xi} f(\xi, \eta),$$

on the other hand if there exists a saddle point  $(\xi_0, \eta_0)$ , then

$$\max_{\xi} \min_{\eta} f(\xi, \eta) \geq \min_{\eta} f(\xi_0, \eta) = f(\xi_0, \eta_0),$$

$$\min_{\eta} \max_{\xi} f(\xi, \eta) \leq \max_{\xi} f(\xi, \eta_0) = f(\xi_0, \eta_0)$$

which gives the other inequality, hence

$$\max_{\xi} \min_{\eta} f(\xi, \eta) = \min_{\eta} \max_{\xi} f(\xi, \eta) = f(\xi_0, \eta_0).$$

With this what needs to be proven is the existence of such a saddlepoint.

<sup>16</sup>For a detailed proof see [von Neumann, 1928, p. 308-309].

<sup>17</sup>"There exists a "saddle point"  $\xi_0, \eta_0$  ( $0 \leq \xi_0 \leq a, 0 \leq \eta_0 \leq b$ ), i.e.,  $f(\xi_0, \eta)$  assumes its minimum for  $\eta = \eta_0$  in  $0 \leq \eta \leq b$  and  $f(\xi, \eta_0)$  assumes its maximum for  $\xi = \xi_0$  in  $0 \leq \xi \leq a$ ." [von Neumann, 1928, (1959 p. 30)].

### The existence of a saddle point

For every fixed  $\xi$  von Neumann considered the set of  $\eta$ ,  $0 \leq \eta \leq b$ , for which  $f(\xi, \eta)$  assumes its minimum value. Since  $f$  is continuous the set will be closed and it will also be convex because  $f$  fullfil the condition (K), this means the set is a subinterval of  $[0, b]$ . Von Neumann let  $[K'(\xi), K''(\xi)]$  denote this subinterval. Thus for fixed  $\xi$ :

$$\{\eta' \in [0, b] \mid \min_{\eta} f(\xi, \eta) = f(\xi, \eta')\} = [K'(\xi), K''(\xi)] \subseteq [0, b].$$

Simillary for fixed  $\eta$ , the set of  $\xi$ ,  $0 \leq \xi \leq a$ , for which  $f(\xi, \eta)$  assumes its maximum, is a closed subinterval of  $[0, a]$ . Von Neumann denoted this subinterval by  $[L'(\eta), L''(\eta)]$ .

That is, for every  $\xi \in [0, a]$  there is an interval  $[K'(\xi), K''(\xi)] \subseteq [0, b]$ , such that every single  $\eta$  belonging to this interval is a point of minimum for the function  $f(\xi, *)$ . Simillary for every  $\eta \in [0, b]$  there is an interval  $[L'(\eta), L''(\eta)] \subseteq [0, a]$ , such that every  $\xi$  in this interval is a point of maximum for the function  $f(*, \eta)$ .

Von Neumann then showed that – due to the continuouity of  $f - K', L'$  and  $K'', L''$  are lower and upper semi-continuous functions respectively. [von Neumann, 1928, p. 310, note 10].

For a fixed  $\xi^*$  von Neumann studied the following set which I have named  $D(\xi^*)$ :

$$D(\xi^*) = \{\xi^{**} \mid \exists \eta^* : \min_{\eta} f(\xi^*, \eta) = f(\xi^*, \eta^*) \text{ and } \max_{\xi} f(\xi, \eta^*) = f(\xi^{**}, \eta^*)\},$$

that is,

$$D(\xi^*) = \cup [L'(\eta^*), L''(\eta^*)] \text{ over } \eta^* \in [K'(\xi^*), K''(\xi^*)].$$

The lower semi-continuous function  $L'$  will assume its minimum value within the interval  $K'(\xi^*) \leq \eta^* \leq K''(\xi^*)$  and the upper semi-continuous function  $L''$  will assume its maximum value. Hence the set  $D(\xi^*)$  will contain a minimal as well as a maximal element. Further more von Neumann argued by means of the following indirect proof that  $D(\xi^*)$  also contain all  $\xi'$  between the minimal and the maximal element: In contradiction to what he wanted to demonstrate von Neumann assumed the existence of an element  $\xi'$  situated in between the minimal and the maximal element but not contained in  $D(\xi^*)$ , then every interval  $[L'(\eta^*), L''(\eta^*)]$  would lie either entirely to the left or entirely to the right of  $\xi'$ . Since  $\xi'$  is between the minimum and the maximum element of  $D(\xi^*)$ , both kind of intervals will exist.  $\eta^*$  runs over an interval,

which implies that both kinds of  $\eta^*$ 's, that is, those  $\eta^*$ 's corresponding to the intervals  $[L'(\eta^*), L''(\eta^*)]$  entirely to the left of  $\xi'$ , and those  $\eta^*$ 's corresponding to the intervals  $[L'(\eta^*), L''(\eta^*)]$  entirely to the right of  $\xi'$ , has a common limit-point  $\eta'$ . This means that both  $L'(\eta^*) \leq \xi'$  and  $L''(\eta^*) \geq \xi'$  will occur arbitrary close to  $\eta'$ , which because of the lower and upper semi-continuity of  $L'$  and  $L''$  respectively, implies that  $L'(\eta') \leq \xi'$  and  $L''(\eta') \geq \xi'$ , that is  $\xi'$  does indeed belong to one of the intervals, namely  $[L'(\eta'), L''(\eta')]$  [von Neumann, 1928, p. 310].

The above result implies that  $D(\xi^*)$  is a closed subinterval of  $[0, a]$ , which von Neumann denoted  $[H'(\xi^*), H''(\xi^*)]$ . To terminate the demonstration von Neumann showed the existence of an element  $\xi^* \in [0, a]$ , which is also a  $\xi^{**}$ , that is, an element  $\xi^*$ , for which  $H'(\xi^*) \leq \xi^* \leq H''(\xi^*)$ . The proof for this is similar to the proof above for the claim that  $D(\xi^*)$  is a closed subinterval, due to the fact that  $H'$  and  $H''$  are lower and upper semi-continuous functions respectively. As before if one assumes that there can exist no such  $\xi^*$  that would imply that all the intervals  $[H'(\xi^*), H''(\xi^*)]$  will lie entirely to the left or entirely to the right of  $\xi^*$ . Again both kinds of  $\xi^{**}$ 's will have a common limit point  $\xi'$  which will belong to the interval  $[H'(\xi'), H''(\xi')]$ .

With this von Neumann has demonstrated the existence of an element  $\xi^* \in [0, a]$ , which also satisfy  $\xi^* \in D(\xi^*)$ . Since this means that there exists an element  $\eta^*$ , such that  $\min_{\eta} f(\xi^*, \eta) = f(\xi^*, \eta^*)$ , and at the same time  $\max_{\xi} f(\xi, \eta^*) = f(\xi^*, \eta^*)$ , the point  $\xi^*, \eta^*$  is a saddle point for the function  $f$ , which finished von Neumann's proof of the minimax theorem.

## 2.4 Von Neumann's 1928 proof in relation to fixed point theorems and systems of inequalities

The 1928-proof of von Neumann is indeed a "tour de force" [Heims, 1980, p. 91]. Regarding the other remarks in the literature that I cited in the introduction it has been said about von Neumann's 1928-proof that he "demonstrates the close connection with fixed-point theorems and especially Brouwer's theorem" [Ingrao and Israel, 1990, p. 211], and that the proof concerns the existence of a solution to a system of equalities and inequalities.<sup>18</sup> These issues do not seem very obvious to me. Von Neumann talked at no point about fixed points and he did not formulate a system of equations and inequalities to be solved.

<sup>18</sup>See [Ingrao and Israel, 1990, p. 211], [Heims, 1980, p. 91].

Yet, in Kuhn and Tucker's paper about von Neumann's work on game theory they wrote:

The analytic proofs of the Minimax Theorem given by von Neumann were of two essentially different types. Proofs of the first type (see [A] and [B]) are based explicitly on extensions of the Brouwers fixed point theorem; [Kuhn and Tucker, 1958, p. 112]

[A] refers to von Neumann's 1928-paper while [B] refers to a paper by von Neumann published in 1937 which will be treated in the next section. In von Neumann's 1928-proof one can "extract" a proof for an extension of Brouwer's fixed point theorem. In doing so the question about existence of a saddle point for the function  $f(\xi, \eta)$  becomes a question about the existence of a fixed point for a 'point to set' map. The connection can be derived in the following way:

In the end von Neumann showed the existence of an element  $\xi^*$  satisfying  $\xi^* \in [H'(\xi^*), H''(\xi^*)]$ . If one puts  $F(\xi^*) = [H'(\xi^*), H''(\xi^*)]$ ,  $F$  can be interpreted as a map for which there to each element  $\xi$  in  $[0, a]$  corresponds a set  $F(\xi)$ , which is a subinterval of  $[0, a]$ . An element  $\xi^*$  which is being mapped onto an interval  $F(\xi^*)$ , which the elements itself belongs to is a kind of a fixed point for a 'point to set' map. With this interpretation the existence of a saddle point and the existence of a fixed point for the mapping  $F$  is one and the same thing.

Von Neumann did not make this interpretation in the 1928-paper and for reasons to be supported in the next section I am not convinced that von Neumann in 1928 was aware of this connection to fixed points.

As far as the connection to systems of linear inequalities and equations is concerned von Neumann in his proof did not draw any connections at all. But as Kuhn and Tucker show in their essay on von Neumann's work it is possible to derive such a connection. It can be done in the following way: In an analysis of the consequences of the minimax theorem for the choice of strategies von Neumann considered the set  $A$  of all  $\xi$  for which  $\min_{\eta} h(\xi, \eta)$  assumes its maximum value  $M$ , and the set  $B$  of all  $\eta$  for which  $\max_{\xi} h(\xi, \eta)$  assumes its minimum value  $M$ . That is,

$$A = \{\xi \in \mathbf{R}^{\Sigma_1} : \min_{\eta} h(\xi, \eta) \text{ assumes its maximum value } M\}$$

$$B = \{\eta \in \mathbf{R}^{\Sigma_2} : \max_{\xi} h(\xi, \eta) \text{ assumes its minimum value } M\}$$

As pointed out by von Neumann it is quite obvious that

1. if  $\xi$  belongs to  $A$  then  $h(\xi, \eta) \geq M$  is always true (because  $h(\xi, \eta) \geq \min_{\eta} h(\xi, \eta) = M$ , since  $\xi$  belongs to  $A$ ),
2. If  $\eta$  belongs to  $B$  then  $h(\xi, \eta) \leq M$  is always true
3. if  $\xi$  does not belong to  $A$  there exists an element  $\eta$  for which  $h(\xi, \eta) < M$ ,
4. if  $\eta$  does not belong to  $B$  there exists an element  $\xi$  for which  $h(\xi, \eta) > M$ ,
5. if  $\xi$  belongs to  $A$  and  $\eta$  belongs to  $B$  then  $h(\xi, \eta) = M$ .

Hence, von Neumann argued, it is obvious that  $S_1$  should choose a strategy  $\xi$  that belongs to  $A$  and  $S_2$  should choose a strategy  $\eta$  which belongs to  $B$ . For every such choice the game will have the value  $M$  for  $S_1$  and the value  $-M$  for  $S_2$  [von Neumann, 1928, p. 305].

In the 1928-paper von Neumann did not discuss this further, but one can make an interpretation of this such that it is concerned with the finding of elements  $\xi^*$ ,  $\eta^*$ , such that the inequalities

$$\xi^* \geq 0, \quad \eta^* \geq 0, \quad \max_{\xi} h(\xi, \eta^*) \leq M, \quad \min_{\eta} h(\xi^*, \eta) \geq M \quad (1)$$

and equalities

$$\xi_1^* + \dots + \xi_{\Sigma_1}^* = 1, \quad \eta_1^* + \dots + \eta_{\Sigma_2}^* = 1 \quad (2)$$

are all satisfied.

Kuhn and Tucker in their essay derived the following connection between solutions  $(\xi^*, \eta^*, M)$  to the 'minimax problem' and a system of linear inequalities and equations. They let  $g(p, q)$  denote the elements of a matrix  $A$ , then  $h(\xi, \eta) = \xi A \eta$ . Hence a solution  $(\xi^*, \eta^*)$  to the linear inequalities and equations

$$\begin{aligned} \xi^* \geq 0, \quad \eta^* \geq 0, \quad A \eta^* \leq M, \quad \xi^* A \geq M \\ \xi_1^* + \dots + \xi_{\Sigma_1}^* = 1, \quad \eta_1^* + \dots + \eta_{\Sigma_2}^* = 1. \end{aligned}$$

will then also be a solution to the system (1) and (2) [Kuhn and Tucker, 1958, p. 111]. But this algebraical interpretation of optimal strategies as constituting a solution to a system of linear equalities and inequalities was

not explicitly formulatet by von Neumann in 1928. As we shall see in the next sections this insight was not to come until later. As a matter of fact von Neumann's 1928 - proof is more general covering also nonlinear functions. To be fair it needs to be said that Kuhn and Tucker did not claim that von Neumann actually made this algebraic characterisation in 1928 but the other statements in the literaure cited above leaves the impression that von Neumann in 1928 was working within a framework of linear inequality theory. As we shall see in the next sections and as von Neumann also himself later remarked to an announcement made by the French mathematician Fréchet, this connection to the theory of convexity and linear inequality theory was only recognized later on.

### 3 The Connection to Fixed Point Theorems and Economy: von Neumann's 1937-Paper

After the 1928-paper 16 years passed by before von Neumann published on game theory again even though the minimax theorem popped up again in 1932 in another disguise though. It happened in a mathematical-economic model that von Neumann developed in the early thirties. The first mention of the work is a talk von Neumann gave on the model at the mathematics seminar at Princeton. The paper was published five years later on a request from Karl Menger under the title "Über ein ökonomisches Gleichungssystem und eine Verallgemeinerung des Brouwerschen Fixpunktsatzes" [von Neumann, 1937].

The model of von Neumann is a linear production model in which he did not distinguish between goods consumed and goods produced in the process of production. He analyzed a situation where there are  $n$  goods  $G_1, \dots, G_n$  which can be produced by  $m$  processes  $P_1, \dots, P_m$ .  $y_1, \dots, y_n$  denotes the prizes of the goods while  $x_1, \dots, x_m$  are the intensities with which the processes are being used. Finally  $a_{ij}$  and  $b_{ij}$  denoted the number of units of the good  $G_j$  consumed and produced respectively by the process  $P_i$ .

Von Neumann was interested in situations where the whole economy expands without change of structure, i.e. where the ratios of the intensities  $x_1 : \dots : x_m$  remain unchanged, although  $x_1, \dots, x_m$  themselves may change [von Neumann, 1937, p. 30]. In such a case the intensities are multiplied by a common factor  $\alpha$  per unit of time, the so-called coefficient of expansion.

The unknowns are the intensities  $x_1, \dots, x_m$ , the coefficient of expansion  $\alpha$ , the prizes  $y_1, \dots, y_n$  of the goods, and the interest factor  $\beta = 1 + \frac{z}{100}$ , where  $z$  is the rate of interest in % per unit of time [von Neumann, 1937, p. 30].

The analysis of von Neumann resulted in the following system of inequalities which were to be solved:

$$x_i \geq 0, \quad (3)$$

$$y_j \geq 0, \quad (4)$$

$$\sum_{i=1}^m x_i > 0, \quad (5)$$

$$\sum_{j=1}^n y_j > 0, \quad (6)$$

$$\alpha \sum_{i=1}^m a_{ij} x_i \leq \sum_{i=1}^m b_{ij} x_i, \quad (7)$$

where  $y_j = 0$  if strict inequality ' $<$ ' holds.

$$\beta \sum_{j=1}^n a_{ij} y_j \geq \sum_{j=1}^n b_{ij} y_j, \quad (8)$$

where  $x_i = 0$  if strict inequality ' $>$ ' holds.

The inequality (7) means that it is impossible to consume more of the good  $G_j$  than the amount produced in the total process. If more is produced than is consumed  $G_j$  becomes a free good with zero prize  $y_j = 0$ . The inequality (8) appears because there is no profit in the model a possible gain would be reinvested. (8) means that in equilibrium there can not be a profit on any process  $P_i$ . If there is a loss, i.e. if ' $>$ ' holds the process  $P_i$  will not be used and  $x_i = 0$  [Von Neumann, 1937, p. 75-76].

### 3.1 The Solution of the System of Inequalities

In order to find necessary and sufficient conditions for the existence of a solution to such a system of linear inequalities von Neumann first transformed

the problem of solutions into a saddle point problem. For this purpose he introduced the function

$$\phi(X, Y) = \frac{\sum_{i=1}^m \sum_{j=1}^n b_{ij} x_i y_j}{\sum_{i=1}^m \sum_{j=1}^n a_{ij} x_i y_j}$$

where  $X = (x_1, \dots, x_m)$  and  $Y = (y_1, \dots, y_n)$  are variables satisfying (3), (5) and (4), (6) respectively. That is, he was looking at the ratio between the total income and the total costs.

Von Neumann then argued that the question of a solution to the system of inequalities (3) – (8) becomes the question of a saddle point for the function  $\phi$ . Hence, he could formulate the question of the existence of a solution to the system (3) – (8) as follows:

(\*) Consider  $(X, Y)$  in the domain bounded by (3) – (6). To find a saddle point  $X = X_0, Y = Y_0$  for  $\phi$ . (See [von Neumann, 1937, p. 78].)

Thus just like in the 1928-paper on games the key problem is to prove the existence of a saddle point for a certain function. Instead of proving the existence of a saddle point right away as he did in the 1928-paper he instead proved a ‘fixed point’-lemma, which is the lemma that appear in the last part of the title: “... eine Verallgemeinerung des Brouwerschen Fixpunktsatzes”. Von Neumann then derived the existence of a saddle point for  $\phi$  as a direct consequence of this lemma. (See [von Neumann, 1937, p. 80].)

### 3.2 The Connection to the Minimax Theorem

At this time von Neumann was fully aware of the connection between the game theoretical problem and the problem of existence of a solution to a system of linear inequalities:

Die Lösbarkeit unseres Problems [The existence of a solution to the system (3) – (8)] hängt sonderbareweise mit jener eines in der Theorie der Gesellschaftsspiele auftretenden Problems zusammen, das der Verf. anderwärtig behandelt hat ... Jenes Problem ist ein Specialfall von (\*) und wird durch unsere Lösung von (\*)

auf eine neue Weise miterledigt.<sup>19</sup> [Von Neumann, 1937, p.79, note 2]

This is the first time von Neumann explicitly states that he has recognized a connection between the solution of systems of linear inequalities and the minimax solution of a two-person zero-sum game. The connection was not trivial. As I pointed out in the previous section, reading von Neumann's 1928-paper on its own terms without making recourse to later developments in game theory there is in my opinion no evidence that von Neumann had realized this connection to systems of linear inequalities in 1928. On the contrary his statement in 1937 that the question about solutions to the system of inequalities is "oddly" connected with the minimax solution shows that this was kind of unexpected. Had he already in 1928 been aware of this he would probably not have called it "odd" ten years later.

Regarding the fixed point technique used by von Neumann to show the existence of a saddle point in the 1937-paper it can be seen both from the title of the paper where the result is announced and from the following quote from the paper that von Neumann found it a quite important result which was interesting in itself

Dieser verallgemeinerte Fixpunktssatz ... ist auch an sich von Interesse.<sup>20</sup> [Von Neumann, 1937, p. 73]

In the previous section I argued that von Neumann in 1928 probably was not aware of the fact that the existence of a saddle point could be proved on the basis of fixed points technique. The above quotation and the fact that he found the generalised fix-point result so important that he announced it in the title indicate that he had not fully recognized this in 1928. If so he would probably had announced it at that time considering the credit he ascribed it in 1937 and there is no mention of fixed points techniques what so ever in the 1928-paper. Another argument in favour of this is that he explicitly wrote in the 1937-paper that the game theoretic problem is solved in the 1937-paper in "a new way".

---

<sup>19</sup> "The question whether our problem has a solution is oddly connected with that of a problem occurring in the Theory of Games dealt with elsewhere. ... The problem there is a special case of (\*) and is solved here in a new way through our solution of (\*)" [von Neumann, 1937, (1945 p. 5, note 1)].

<sup>20</sup> "This generalised fix-point theorem ... is also interesting in itself." [von Neumann, 1937 (1945 p. 1)].

## 4 The Minimax Theorem in the Theory of Convexity: The 1944–Proof

In 1937 von Neumann was as we have just seen aware of the connection between the minimax theorem and solutions of systems of linear inequalities. The proof though was not build on the algebra of inequalities but was founded on topological methods. The first algebraic proof of the minimax theorem was due to the French mathematician Jean Ville who published it in Émile Borel's book "Traité du calcul des probabilités et de ses applications" from 1938.

Borel himself had published a series of notes on games from 1921 to 1927. He was the first one who tried to build a mathematical theory for games but after the publication of von Neumann's minimax theorem in 1928 he seemed to have lost interest in the subject<sup>21</sup>. He published a note on game theory in 1927 and then he did not publish anything on games until this book of probability came out 10 years later. In the book Borel has a chapter written by himself devoted to game theory and quite strickingly there is no reference at all to von Neumann and the minimax theorem in that chapter.<sup>22</sup> Instead the minimax theorem is treated in a separate note by Jean Ville with the title "Théorème de M. von Neumann" [Ville, 1938].

Ville's algebraic proof is important because it exercised a direct influence on von Neumann during von Neumann's work with the first collected and coherent book on game theory and thereby gave rise to a development which led to the establishment of the minimax theorem in the theory of convexity.

I will only present the key tools in Ville's proof and not go into detail with the proof itself.<sup>23</sup> Ville derived his key tool as a corollar to the following lemma concerning linear forms, which he proved by induction:

Let  $p$  linear forms in  $n$  variables be given:

$$f_j(x) = \sum a_j^i x_i \quad (j = 1, \dots, p; i = 1, \dots, n).$$

---

<sup>21</sup>Before the work of Borel one only finds attempts to mathematize specific games like the card game "le Her" by James Waldegrave, baccarat by Joseph Bertrand in 1899 and chess by Ernst Zermelo in 1913. For accounts on these earlier attempts and on the work of Borel see [Dimand and Dimand, 1992]. For accounts on the work of Borel see also [Leonard, 1992].

<sup>22</sup>In [Leonard, 1992] Leonard discuss' this issue and concludes that "This can only be regarded as an act of deliberate omission by Borel." [Leonard, 1992, p. 46].

<sup>23</sup>See also [Leonard, 1992]. For further details on the proof see [Kjeldsen, 1999].

Suppose they have the following property:<sup>24</sup>

For all  $x \geq 0$  there exists a  $j$  in  $\{1, \dots, p\}$ , such that  $f_j(x) \geq 0$ .

Then the following hold true: There exists at least one set of nonnegative coefficients

$$X_1, \dots, X_p \text{ with } X_1 + \dots + X_p = 1,$$

such that

$$\sum_{j=1}^p X_j f_j(x) \geq 0 \text{ for all } x \geq 0.$$

[Ville, 1938, p. 105]

The key tool Ville was able to derive from this lemma was the following result

Let  $f_1, \dots, f_p$  be  $p$  linear forms in  $n$  variables  $x_1, \dots, x_n$ , and let  $\phi$  be a linear form in the same variables. If for every point  $x \geq 0$  at least one of the forms  $f_j$  assumes a value greater than or equal to the value of  $\phi$  then a linear combination

$$\psi = X_1 f_1 + \dots + X_p f_p, \quad X_j \geq 0, \quad X_1 + \dots + X_p = 1,$$

exists for which  $\psi \geq \phi$  for all  $x \geq 0$ . [Ville, 1938, p. 107]

From this result Ville gave a fairly easy proof of von Neumann's minimax theorem.

#### 4.1 Von Neumann's 1944-proof

The final placement of game theory in general and the minimax theorem in particular within a context of linear inequalities and the theory of convexity was due to the joint work "Theory of Games and Economic Behavior" by von Neumann and the Austrian economist Oskar Morgenstern [von Neumann and Morgenstern, 1944]. According to Kuhn and Tucker the proof of the minimax theorem which von Neumann and Morgenstern presented in their book was inspired directly by the proof given by Ville:

<sup>24</sup> $x \geq 0$  means  $x_i \geq 0$  for every  $i = 1, \dots, n$ .

Oskar Morgenstern has told us [Kuhn and Tucker] that he drew Ville's article to von Neumann's attention after seeing it quite by chance while browsing in the library of the Institute for Advanced Study. They decided at once to adopt a similar elementary procedure, trying to make it as pictorial and simple to grasp as possible. [Kuhn and Tucker, 1958, p. 116]

How thrilled Morgenstern was when he discovered Ville's proof is evident from a note in his diary dated christmas eve 1941:

Both [the 1938-book of Borel and the proof by Ville] are unknown to Johnny. Now he has discovered additional proofs that are becoming increasingly simple and are purely algebraic!! It necessitates some modification in the text, but we can print it. (Quoted in [Rellstab, 1992, p. 87])

The new proofs by von Neumann that Morgenstern speaks about were indeed very different from the earlier proofs by von Neumann. The proof they gave in "Theory of Games and Economic Behavior" is, as we shall see, of a purely algebraic nature and falls within what von Neumann and Morgenstern themselves characterised as

the mathematico-geometrical theory of linearity and convexity. [von Neumann and Morgenstern, 1944, p. 128]

### **The Theorem of the Alternative for Matrices**

The essential tool in the proof of the minimax theorem that Morgenstern and von Neumann gave in 1944 is what they called "The Theorem of the Alternative for Matrices". They derived this theorem as a direct consequence of the theorem of supporting hyperplanes which states that

Given  $x_1, \dots, x_p$  in  $\mathbf{R}^n$ . Then a  $y$  in  $\mathbf{R}^n$  either belongs to the convex set  $C$  spanned by  $x_1, \dots, x_p$ , or there exists a hyperplane which contain  $y$  such that  $C$  falls entirely within one half-space produced by that hyperplane. [von Neumann and Morgenstern, 1944, p. 134]

Von Neumann and Morgenstern considered a  $(n \times m)$  matrix, let us call it  $A$ , with elements  $a(i, j)$ ,  $i = 1, \dots, n$ ;  $j = 1, \dots, m$ . In order to use the

theorem of supporting hyperplanes von Neumann and Morgenstern formed the convex set  $C$  spanned by the  $m$  column vectors in  $A$  together with the  $n$  coordinate vectors in  $\mathbf{R}^n$ . Putting  $y = 0$  either  $0$  belongs to  $C$  or to a hyperplane  $H$ , such that all of  $C$  is contained in one half-space produced by that hyperplane [von Neumann and Morgenstern, 1944, p. 139]. In the first case they could prove the existence of  $x$  in  $\mathbf{R}^m$  for which  $x_1 \geq 0, \dots, x_m \geq 0$ ,  $\sum_{j=1}^m x_j = 1$ , such that the inequalities

$$\sum_{j=1}^m a(i, j)x_j \leq 0$$

are satisfied for  $i = 1, \dots, n$ . In the second case, that is where  $0$  does not belong to  $C$ , they showed the existence of a vector  $w$  in  $\mathbf{R}^n$  with  $w_1 > 0, \dots, w_n > 0$ ,  $\sum_{i=1}^n w_i = 1$ , such that the following inequalities are satisfied:

$$\sum_{i=1}^n a(i, j)w_i > 0 \quad \text{for } j = 1, \dots, m.$$

These two possibilities or alternatives as von Neumann and Morgenstern called them exclude each other. Formulated in matrix notation they had proved the following result:

If  $A$  is a  $(n \times m)$  matrix then exactly one of the following two systems of inequalities has a solution:

$$Ax \leq 0, \quad x \geq 0, \quad \sum_{j=1}^m x_j = 1,$$

$$wA > 0, \quad w > 0, \quad \sum_{i=1}^n w_i = 1.$$

[von Neumann and Morgenstern, 1944, p. 138-141]

From this result they proved the minimax theorem for two-person zero-sum games in the following way: Keeping the notation from von Neumann's 1928-paper they let

$$h(\xi, \eta) = \sum_{p=1}^{\Sigma_1} \sum_{q=1}^{\Sigma_2} g(p, q)\xi_p\eta_q$$

be the expected value for the player  $S_1$ . By defining  $A$  to be the matrix  $(g(p, q))_{(\Sigma_1 \times \Sigma_2)}$ , they obtained either the existence of a vector  $\xi \in \mathbf{R}^{\Sigma_1}$  for which  $\xi \geq 0$ ,  $\sum \xi_p = 1$ , such that

$$\sum_{p=1}^{\Sigma_1} g(p, q) \xi_p \geq 0 \quad \text{for } q = 1, \dots, \Sigma_2, \quad (9)$$

or the existence of a vector  $\eta \in \mathbf{R}^{\Sigma_2}$  for which  $\eta \geq 0$ ,  $\sum \eta_q = 1$ , such that

$$\sum_{q=1}^{\Sigma_2} g(p, q) \eta_q \leq 0 \quad \text{for } p = 1, \dots, \Sigma_1. \quad (10)$$

If (9) holds true then

$$v_1 = \max_{\xi} \min_{\eta} h(\xi, \eta) \geq 0.$$

If on the other hand (10) holds true then

$$v_2 = \min_{\eta} \max_{\xi} h(\xi, \eta) \leq 0.$$

From this von Neumann and Morgenstern concluded that

$$\text{either } v_1 \geq 0 \quad \text{or} \quad v_2 \leq 0$$

that is *never*

$$v_1 < 0 < v_2. \quad (11)$$

The final step in the proof was to show that (11) never can be the case not only for 0 but for an arbitrary number  $w$ , that is *never*

$$v_1 < w < v_2.$$

Since  $v_1 \leq v_2$  is always true von Neumann and Morgenstern had proven the equality:

$$v_1 = \max_{\xi} \min_{\eta} h(\xi, \eta) = \min_{\eta} \max_{\xi} h(\xi, \eta) = v_2.$$

In this way von Neumann and Morgenstern in 1944 reduced the proof for the minimax theorem to a fairly simple consequence of the theorem of "Alternatives for Matrices" which is a purely algebraic theorem with in the theory of systems of linear inequalities.

## 5 Conclusion on von Neumann's Perception of the Minimax Theorem

The above analysis of the development of von Neumann's understanding of the different mathematical contexts the minimax theorem presented itself in during the period 1928 to 1944 clearly shows that his recognition of the connections between the minimax theorem on the one hand and fix-point results and the theory of linear inequalities on the other hand only emerged gradually. The full understanding of the connection to fixed points theorems was not present until 1937 while the final establishment and realization of the minimax theorem as a result belonging to the theory of linear inequalities and the theory of convexity was not fully recognized until 1944.

It can also serve as an illustration of how mathematics evolves. In this case a problem (the solution of a two-person zero-sum game) emerged in connection with a new kind of mathematical questions (the mathematization of games). The problem is solved in itself and in the beginning possible connections to other branches of mathematics can be very difficult sometimes impossible to realize. Later the problem or a similar problem crops up again in another context (the economic model of von Neumann) one recognize the connection and simultaneously the complexity of the problem decomposes, the underlying structure of the proof becomes visible (fixed points techniques) and new general results (the extension of Brouwer's fixed point theorem) which are interesting in themselves not limited to the context they originally were derived in emerges. Finally the problem is recognized to be a simple consequence of fundamental theorems in a different branch of mathematics (theories of linear inequality and convexity).

## 6 The Discussion of Priority: the Significance of the Minimax Theorem

The 1928-paper by von Neumann was generally known to mark the beginning of game theory. The earlier notes on the subject from the beginning of the twenties by Borel was not generally known until after 1953 when the French mathematician Maurice Fréchet had three of them translated into English. In the introduction to the translation Fréchet argued for the importance of Borel's work:

It was only relatively recently that I began to occupy myself with the theory of probability and its applications, which explains why the notes that Émile Borel . . . published between 1921 and 1927 on the theory of psychological games escaped my attention. It was chance to begin with . . . because, in the extensive literature devoted to this theory [game theory] and its applications in recent years, references to earlier work do not lead back, in general, further than to the important paper published in 1928 by Professor von Neumann. But, in reading these notes of Borel's I discovered that in this domain, as in so many others, Borel had been an initiator. [Fréchet, 1953a, p. 95]

In order to understand the priority debate and how it connects to the significance of the context to be discussed below, we need to know a little about Borel's work.

## 6.1 Borel's work on game theory

The first of Borel's notes on the subject was published in 1921 [Borel, 1921]. He considered a symmetric game with two players  $A$  and  $B$ . He introduced the concept "méthode de jeu" (method of play) and the fundamental question asked by Borel was then, whether it was possible to determine a "méthode de jeu meilleure" (Best method of play). It was not quite clear what was to be understood by a "méthode meilleure" but his concept of "méthode de jeu" was the same as von Neumann's, that is what now is called a pure strategy [Borel, 1921, p. 1304]. Like von Neumann Borel assumed the players had a finite number of strategies  $C_1, \dots, C_n$  to choose from.

Borel's inspiration to investigate games came from his work on probability and in his first paper he was looking for the probabilities for winning the game. His starting point was, that if  $A$  chooses the strategy  $C_i$ , and  $B$  chooses the strategy  $C_k$ , then the probability that  $A$  wins the game can be calculated. He called that probability for  $a$ . The probability for player  $B$  is then  $b = 1 - a$ . To indicate that these probabilities are dependent of the choices of strategies he put

$$a = \frac{1}{2} + \alpha_{ik}$$

and

$$b = \frac{1}{2} + \alpha_{ki}$$

where  $\alpha_{ik}$  and  $\alpha_{ki}$  lie between  $-\frac{1}{2}$  and  $+\frac{1}{2}$  and satisfy the relation

$$\alpha_{ik} + \alpha_{ki} = 0.$$

Like von Neumann he also considered the concept of what later became known as mixed strategies [Borel, 1921, p. 1305]. But in contrary to von Neumann who considered the actions of both players simultaneously Borel began by examine singular cases calculating if it would be possible for one of the players to choose a mixed strategy such that the probability that he or she would win would be  $\frac{1}{2}$  no matter what strategy the other player would choose. In the 1921-note he calculated for the case where there are only three pure strategies to choose from and he reached a possitive conclusion. In general though he was convinced that for games with more than three pure strategies the answer would be negative [Borel, 1921, p. 1306]. Two years later he had done the calculations for games with five pure strategies which shows that the answer also in this case turned out to be positive and he thought that it would probably also be true for seven pure strategies, but he still thought that for a larger number of strategies the answer should be no [Borel, 1923, p. 1117].

In 1924 Borel included a chapter on games in his book on probability [Borel, 1924]. Instead of looking at the probabilities for winning as he did in 1921, he now let  $\alpha_{ik}$  denote an amount of money, which player  $B$  has to give to player  $A$ , if player  $A$  chooses strategy  $C_i$  and player  $B$  chooses strategy  $C_k$ . The question he was trying to answer then was, is it possible for player  $A$  to choose a mixed strategy such that the expected value he or she can get is 0, no matter which mixed strategy player  $B$  chooses? That is, can player  $A$  choose a strategy that in all cases can protect  $A$  from loosing money? This is in principle the same question that lead von Neumann to the minimax theorem namely, what is the best you can do in the worst possible case, which is the case where your opponent some how has gained knowledge about your choice of strategy.

Changing the interpretation of the  $\alpha_{ik}$ 's did not make Borel to change his mind about the answer to the question under consideration. He still believed that if the number of pure strategies were larger than seven, the answer would be no. Two years later he had not yet found an argument for his believe and in a note from 1926 he formulated both situations e.i., for a symmetric two-person game is it allways possible for player  $A$  to chose a mixed strategy for which the expected value of the game will be 0 no matter what strategy

player *B* choses or is it not the case? [Borel, 1926]. The second situation contradicts the minimax theorem. The fact that he formulated the positive situation first has been interpreted as he seriously doubted his original views and was beginning to believe that maybe what is now called the minimax theorem would turn out to be true.<sup>25</sup>

The reason why Borel at the outset did not believe in a positive answer has been discussed by Luca Dell'Aglio in the paper "Divergences in the History of Mathematics: Borel, von Neumann and the Genesis of Game Theory" [Dell'Aglio, 1995]. Dell'Aglio argues that Borel had a psychological interpretation of the concept of mixed strategies which

... constitute the conceptual basis of Borel's negation of the minimax theorem in his earlier research into game theory. [Dell'Aglio, 1995, p. 21]

The psychological interpretation enters the picture because Borel on several occasions talks about the advantage of being a better psychologist. The player who is a better observant and analyst than the opponent will have an advantage in the game, which is not true for optimal solutions covered by the minimax theorem. Dell'Aglio concludes that

... the divergence over the validity of the minimax theorem was ultimately due to a difference in the conceptual and technical structure underlying the two theories. In other words, Borel and von Neumann produced different theoretical forecasts because they were working on different basic problems. [Dell'Aglio, 1995, p. 40]

The two different problems that Dell'Aglio is referring to, emerges because von Neumann's point of departure was "the possibility of the existence of equilibria in games played by equal players" [Dell'Aglio, 1995, p. 40] while "Borel took into consideration a similar problem but supposing one player has acquainted himself with the psychological characteristics of his opponent" [Dell'Aglio, 1995, p. 40].

I do not quite understand in what sense Dell'Aglio mean that von Neumann and Borel are studying two different problems. Both of them had as point of departure that you choose your strategy without knowing what your opponent are going to choose. In Borel's calculations for the cases with three

---

<sup>25</sup>See e.g. [Fréchet, 1953b, p. 122].

and five pure strategies he are, like von Neumann, looking for a strategy that can protect you from being in a losing position, no matter what the other player does. That is, a strategy where the result of the game will not change to your disadvantage even though your opponent somehow found out which one you picked, and in a situation like that it does not matter which one is a better psychologist. As far as I can see the main difference between their work is the approach in their investigations. Borel did not consider the mixed strategies of the two players at the same time. He did not work on the interplay of the two players simultaneously and independent choice of strategy. Von Neumann did so and that brought the various 'minmax' and 'maxmin' considerations into the picture and it is precisely the interaction between those that made him realize the solution as a saddle point.

## 6.2 Discussion of Priority

In the quotation previously cited from the introduction by Fréchet to the translation of the notes of Borel, Fréchet announced Borel to be an "initiator" in the domain of game theory. In a commentary Fréchet argued for this opinion:

Borel was the first to indicate the potential importance for this theory of knowing whether this theorem [the minimax theorem], applied to  $n$  manners of playing, is true for arbitrary  $n$ . He did, moreover, demonstrate it for  $n = 3$  and  $n = 5$ , but only for these values. [Fréchet, 1953b, p. 122]

This introduction and commentary of Fréchet caused a brief priority discussion between von Neumann and Fréchet. According to L. J. Savage, the translator of Borel's work, von Neumann got very angry when he learned what Fréchet had written [Heims, 1980, p. 440, note 14]. Von Neumann acknowledged that Borel had been the first one to introduce the concepts of pure and mixed strategies but, he continued,

The relevance of this concept [of mixed strategies] in his [Borel's] hands was essentially reduced by his failure to prove the decisive 'minimax theorem', or even to surmise its correctness. As far as I can see, there could be no theory of games on these bases without that theorem. ... I felt that there was nothing worth publishing until the 'minimax theorem' was proved. [von Neumann, 1953, p. 124-125]

What I find interesting is not so much the priority debate in itself but more the following remark by Fréchet which shows a view very different from that of von Neumann on the importance of the minimax theorem:

Again, it may be mentioned, that even if Borel had, before von Neumann, established the minimax theorem in its full generality; the profound originality of Borel's notes would not have been augmented nor even touched from the economic point of view. He would not thereby have even enriched the set of properly mathematical discoveries for which Borel has acquired a world-wide reputation. He would have, like von Neumann, simply entered an open door. . . . the samme theorem and even more general theorems had been independently demonstrated by several authors well before the notes of Borel and the first paper of von Neumann. [Fréchet, 1953b, p. 122]

The proofs Fréchet is referring to are proofs of theorems similar to von Neumann and Morgenstern's "Alternatives for Matrices", e.i., theorems about solutions to systems of linear inequalities by Minkowski, Farkas, Stiemke, and Weyl.<sup>26</sup>

In 1953 the minimax theorem was realized to be a simple consequence of those classical theorems about solutions to systems of linear inequalities, but von Neumann derived the minimax theorem in a theory of "Gesellschaftsspiele" which was a completely different mathematical context. The techniques used by von Neumann in 1928 had at first nothing to do with linear inequalities, it was not until Ville's proof in 1938 that this connection was recognized, a connection von Neumann and Morgenstern then developed further in their 1944-book. But as von Neumann's 1928 - proof and his 1937 - proof clearly demonstrate and as he himself wrote in 1953 in his answer to Fréchet:

This connection may now seem very obvious to someone who first saw the theory after it had obtained its present form. (O. Morgenstern and myself, in our presentation in 1943, made, for didactical reasons, every effort to emphasize this connection.) However, this was not at all the aspect of the matter in 1921 - 1938. The theorem, and its relation to the theory of convex sets were far from

---

<sup>26</sup>See [Farkas, 1901, p. 5-7], [Stiemke, 1915, p. 340], [Gordan, 1873, p. 23-28], [Minkowski, 1896, p. 39-45], [Weyl, 1935].

being obvious . . . . It is common and tempting fallacy to view the later steps in a mathematical evolution as much more obvious and cogent after the fact than they were beforehand. [von Neumann, 1953, p. 125]

## 7 Conclusion on the Significance of the Context

In this discussion Fréchet advocates for the point of view, that the significance of a mathematical theorem is independent of the mathematical context in which it was derived. The history of von Neumann's development and conception of the minimax theorem shows, that it was far from being trivial and took a larger effort to realize the connection between solutions of systems of linear inequalities and the existence of optimal strategies for two-person zero-sum games. The fact, that the minimax theorem later turned out to be a simple consequence of theorems of inequalities proved earlier, does not render the minimax theorem superfluous or worthless in relation to the development of mathematics, as Fréchet seems to imply. In his assessment whether the minimax theorem has "enriched the set of properly mathematical discoveries" or not, an assessment of the significance of the theorem for the developing of new mathematics is lacking. The mathematical context in which a result is derived determines its formulation and interpretation, and thereby also to which kind of new research it can lead. The questions that guide the research in game theory are not necessarily the same as those guiding the research in the abstract theory of linear inequalities. Hence, the minimax theorem can from a game theoretic point of view be very different from the theorems of linear inequalities.

The minimax theorem of von Neumann had a tremendous influence on the further development of game theory which became a very active field of research after World War II.<sup>27</sup> It also had a decisive influence on the development of some new disciplines in applied mathematics especially linear and nonlinear programming which originated in connection with the Second World War.<sup>28</sup>

---

<sup>27</sup>See for example [Mirowski, 1991].

<sup>28</sup>See [Kjeldsen, 1999, 2000].

## References

- Borel, É. (1921): "La théorie du jeu et les équations intégrales à noyau symétrique." *Comptes Rendus de l'Académie des Sciences*, dec. 19, 173, 1921, pp. 1304-1308.
- Borel, É. (1923): "Sur les jeux où le hasard se combine avec l'habileté des joueurs." *Comptes Rendus de l'Académie des Sciences*, dec. 26, 1923, pp. 1117-1118.
- Borel, É. (1924): "Sur les jeux où interviennent le hasard et l'habileté des joueurs." in É. Borel: *Théorie des Probabilités*. Paris: Librairie Scientifique, J.Hermann, 1924, pp. 204-224.
- Borel, É. (1926): "Un théorème sur les systèmes de formes linéaires à déterminant symétrique gauche." *Comptes Rendus de l'Académie des Sciences*, 183, nov. 22, 1926, pp. 925-927.
- Borel, É. et al. (1938): *Traité du Calcul des probabilités et de ses applications*. Paris: Gauthier-Villars, 1938.
- Dell'Aglio, L. (1995): "Divergences in the History of Mathematics: Borel, von Neumann and the Genesis of Game Theory." *Rivista di Storia della Scienza*, 3, 1995, pp. 1-46.
- Dimand, R. W. and Dimand, M. A. (1992): "The Early History of the Theory of Strategic Games from Waldegrave to Borel." in E. R. Weintraub (ed.) *Towards a History of Game Theory*. Durham and London: Duke University Press, 1992, pp. 15-28.
- Farkas, J. (1901): "Theorie der einfachen Ungleichungen." *Journal für die reine und angewandte Mathematik*, 124, 1901, pp. 1-27.
- Fréchet, M. (1953a): "Emile Borel, Initiator of the Theory of Psychological Games and its Application." *Econometrica*, 21, 1953, pp. 95-96.
- Fréchet, M. (1953b): "Commentary on the three notes of Emile Borel." *Econometrica*, 21, 1953, pp. 118-124.
- Gordan, P. (1873): "Über die Auflösungen linearer Gleichungen mit reellen Coefficienten." *Mathematische Annalen*, 6, 1873, pp. 23-28.

- Heims, S. J. (1980): *John von Neumann and Norbert Wiener*. Cambridge, Massachusetts: The MIT Press, 1980.
- Ingraro, B. and Israel, G. (1990): *The Invisible Hand*. Cambridge, Massachusetts: The MIT Press, 1990.
- Kjeldsen, T. H. (1999b): "A Contextualised Mathematico-historical Analysis of Nonlinear Programming: Development and Multiple Discovery." (in Danish), IMFUFA, Text 372, Roskilde University, 1999.
- Kjeldsen, T. H. (2000): "A Contextualized Historical Analysis of the Kuhn-Tucker Theorem in Nonlinear Programming: The Impact of World War II." Accepted for publication in *Historia Mathematica*, August, 2000.
- Kuhn, H. W. and Tucker, A. W. (1958): "John von Neumann's Work in the Theory of Games and Mathematical Economics." *Bulletin of the American Mathematical Society*, 64, 1958, pp. 100-122.
- Leonard, R. J. (1992): "Creating a Context for Game Theory." in E. Roy Weintraub (ed.): *Towards a History of Game Theory*. Durham and London: Duke University Press, 1992, pp. 29-76.
- Leonard, R. J. (1995): "From Parlor Games to Social Science: von Neumann, Morgenstern, and the Creation of Game Theory 1928-1944." *The Journal of Economic Literature*, 33, 1995, pp. 730-761.
- Minkowski, H. (1896): *Geometrie der Zahlen*. Leipzig: B. G. Teubner, 1896, pp. 39-45.
- Mirowski, P. (1991): "When Games Grow Deadly Serious: The Military Influence on the Evolution of Game Theory." in D. G. Goodwin (ed.): *Economics and National Security*, Annual Supplement to Volume 23, History of Political Economy. Durham and London: Duke University Press, 1991, pp. 227-255.
- Mirowski, P. (1992): "What Were von Neumann and Morgenstern Trying to Accomplish?" in E. Roy Weintraub (ed.): *Toward a History of Game Theory*. Durham and London: Duke University Press, 1992, pp. 113-147.
- Rellstab, U. (1992): "New Insights into the Collaboration between John von Neumann and Oskar Morgenstern on the *Theory of Games and Economic*

- Behavior.*" in E. Roy Weintraub (ed.): *Towards a History of Game Theory*. Durham and London: Duke University Press, 1992, pp. 77-94.
- Stiemke, E. (1915): "Über positive Lösungen homogenen linearer Gleichungen." *Mathematische Annalen*, 76, 1915, pp. 340-342.
- Ulam, S. (1958): "John von Neumann, 1903-1957." *Bulletin of the American Mathematical Society*, 64, 1958, pp. 1-49.
- Ville, J. (1938): *Sur la Théorie générale des jeux où intervient l'habileté des joueurs*. in É. Borel et al. (eds.): *Traité du Calcul des probabilités et de ses applications*. Paris: Gauthier-Villars, 1938, pp. 105-117.
- von Neumann, J. (1928a): "Sur la théorie des jeux." *Comptes Rendus de l'Académie des Sciences*, 186.25 (18. of June), 1928, pp. 1689-1691.
- von Neumann, J. (1928): "Zur Theorie der Gesellschaftsspiele." *Mathematische Annalen*, 100, 1928, pp. 295-320.
- von Neumann, J. (1937): "Über ein ökonomische Gleichungssystem und eine Verallgemeinerung des Brouwerschen Fixpunktsatzes." in K. Menger (ed.) *Ergebnisse eines Mathematischen Kolloquiums*. Wien, 1937, pp. 73-83.
- von Neumann, J.: "A Model of General Economic Equilibrium." (English translation of [von Neumann, 1937]) in *Review of Economic studies*, vol. 13, 1945, pp. 1-9.
- von Neumann, J. (1953): "Communication on the Borel Notes." *Econometrica*, 21, 1953, pp. 124-125.
- von Neumann, J. (1959): "On the Theory of Games of Strategy." (English translation of [von Neumann, 1928]) in A. W. Tucker and R. D. Luce (eds.): *Contributions to the Theory of Games*. Princeton, New Jersey: Princeton University Press, 1959, pp. 13-42.
- von Neumann, J. (1963): *Collected Works*, (ed.:) A. H. Taub, Pergamon Press: New York, 1963.
- von Neumann, J. and Morgenstern, O. (1944): *Theory of Games and Economic Behavior*. Princeton: Princeton University Press, Princeton, 1944.

Weyl, H. (1935): "Elementare Theorie der konvexen Polyeder." *Mathematici Helvetici*, 7, 1935, pp. 209-306.

Liste over tidligere udsendte tekster kan rekvireres  
på IMFUFA's sekretariat, tlf. 4674 2263 eller  
e-mail: bs@ruc.dk

- 332/97 ANOMAL SWELLING AF LIPIDE DOBBELTLAG  
Specialerapport af:  
Stine Sofia Korremann  
Vejleder: Dorthe Posselt
- 333/97 Biodiversity Matters  
an extension of methods found in the literature  
on monetisation of biodiversity  
by: Bernd Kuemmel
- 334/97 LIFE-CYCLE ANALYSIS OF THE TOTAL DANISH  
ENERGY SYSTEM  
by: Bernd Kuemmel and Bent Sørensen
- 335/97 Dynamics of Amorphous Solids and Viscous Liquids  
by: Jeppe C. Dyre
- 336/97 PROBLEM-ORIENTATED GROUP PROJECT WORK AT  
ROSKILDE UNIVERSITY  
by: Kathrine Legge
- 337/97 Verdensbankens globale befolkningsprognose  
- et projekt om matematisk modellering  
af: Jørn Chr. Bendtsen, Kurt Jensen,  
Per Pauli Petersen  
Vejleder: Jørgen Larsen
- 338/97 Kvantisering af nanolederes elektriske  
ledningsevne  
Første modul fysikprojekt  
af: Søren Dam, Esben Danielsen, Martin Niss,  
Esben Friis Pedersen, Frederik Resen Steenstrup  
Vejleder: Tage Christensen
- 339/97 Defining Discipline  
by: Wolfgang Coy
- 340/97 Prime ends revisited - a geometric point  
of view -  
by: Carsten Lunde Petersen
- 341/97 Two chapters on the teaching, learning and  
assessment of geometry  
by Mogens Niss
- 342/97 LONG-TERM SCENARIOS FOR GLOBAL ENERGY  
DEMAND AND SUPPLY  
A global clean fossil scenario discussion paper  
prepared by Bernd Kuemmel  
Project leader: Bent Sørensen
- 343/97 IMPORT/EKSPORT-POLITIK SOM REDSKAB TIL OPTIMERET  
UDNYTTELSE AF EL PRODUCERET PÅ VE-ANLÆG  
af: Peter Meibom, Torben Svendsen, Bent Sørensen
- 344/97 Puzzles and Siegel disks  
by Carsten Lunde Petersen
- 
- 345/98 Modeling the Arterial System with Reference to  
an Anesthesia Simulator  
Ph.D. Thesis  
by: Mette Sofie Olufsen
- 346/98 Klyngedannelse i en hulkatode-forstøvningsproces  
af: Sebastian Horst  
Vejledere: Jørn Borggren, NBI, Niels Boye Olsen
- 347/98 Verificering af Matematiske Modeller  
- en analyse af Den Danske Eulerske Model  
af: Jonas Blomqvist, Tom Pedersen, Karen Timmermann.  
Lisbet Øhlenschläger  
Vejleder: Bernhelm Booss-Bavnbek
- 348/98 Case study of the environmental permission  
procedure and the environmental impact assessment  
for power plants in Denmark  
by: Stefan Krüger Nielsen  
Project leader: Bent Sørensen
- 349/98 Tre rapporter fra FAGMAT - et projekt om tal  
og faglig matematik i arbejdsmarkedsuddannelserne  
af: Lena Lindenskov og Tine Wedege
- 350/98 OPGAVERSAMLING - Bredde-Kursus i Fysik 1976 - 1998  
Erstatter teksterne 3/78, 261/93 og 322/96
- 351/98 Aspects of the Nature and State of Research in  
Mathematics Education  
by: Mogens Niss

- 352/98 The Herman-Swiatec Theorem with applications  
by: Carsten Lunde Petersen
- 353/98 Problemløsning og modellering i en almindelig matematikundervisning  
Specialerapport af: Per Gregersen og Tomas Højgaard Jensen  
Vejleder: Morten Blomhøj
- 354/98 A GLOBAL RENEWABLE ENERGY SCENARIO  
by: Bent Sørensen and Peter Meibom
- 355/98 Convergence of rational rays in parameter spaces  
by: Carsten Lunde Petersen and Gustav Ryd
- 356/98 Terrænmodellering  
Analyse af en matematisk model til konstruktion af terrænmodeller  
Modelprojekt af: Thomas Frommelt, Hans Ravnkjær Larsen og Arnold Skimminge  
Vejleder: Johnny Ottesen
- 357/98 *Cayleys Problem*  
En historisk analyse af arbejdet med Cayley problem fra 1870 til 1918  
Et matematisk videnskabsfagsprojekt af: Rikke Degn, Bo Jakobsen, Bjarke K.W. Hansen, Jesper S. Hansen, Jesper Udesen, Peter C. Wulff  
Vejleder: Jesper Larsen
- 358/98 *Modeling of Feedback Mechanisms which Control the Heart Function in a View to an Implementation in Cardiovascular Models*  
Ph.D. Thesis by: Michael Danielsen
- 359/99 *Long-Term Scenarios for Global Energy Demand and Supply Four Global Greenhouse Mitigation Scenarios*  
by: Bent Sørensen
- 360/99 **SYMMETRI I FYSIK**  
En Meta-projektrapport af: Martin Niss, Bo Jakobsen & Tune Bjarke Bonnê  
Vejleder: Peder Voetmann Christiansen
- 361/99 *Symplectic Functional Analysis and Spectral Invariants*  
by: Bernhelm Booss-Bavnbek, Kenro Furutani
- 362/99 *Er matematik en naturvidenskab? - en udspænding af diskussionen*  
En videnskabsfagsprojekt-rapport af Martin Niss  
Vejleder: Mogens Nørgaard Olesen
- 363/99 **EMERGENCE AND DOWNWARD CAUSATION**  
by: Donald T. Campbell, Mark B. Bickhard and Peder V. Christiansen
- 364/99 *Illustrationens kraft*  
Visuel formidling af fysik  
Integreret speciale i fysik og kommunikation af: Sebastian Horst  
Vejledere: Karin Beyer, Søren Kjørup
- 365/99 *To know - or not to know - mathematics, that is a question of context*  
by: Tine Wedege
- 366/99 **LATEX FOR FORFATTERE**  
En introduktion til LATEX og IMFUPA-LATEX af: Jørgen Larsen
- 367/99 **Boundary Reduction of Spectral invariants and Unique Continuation Property**  
by Bernhelm Booss-Bavnbek
- 368/99 Kvartalsrapport for projektet  
Scenarier for samlet udnyttelse af brint som energibærer i Danmarks fremtidige energisystem  
Projektleder: Bent Sørensen  
Opdateret til halvvejsrapport. Den nye udgave Tekst 368bis kan hentes ned fra internettet fra adressen <http://mmf.ruc.dk/energy/report>
- 369/99 Dynamics of Complex Quadratic Correspondences  
by: Jacob Jalving
- 370/99 **OPGAVESAMLING**  
Bredde-Kursus i Fysik 1976 - 1999 (erstatte tekst nr. 350/98)
- 371/99 Bevisets stilling - beviser og bevisførelse i en gymnasial matematikundervisning  
Matematikspeciale af: Maria Hermansson  
Vejleder: Mogens Niss
- 372/99 En kontekstualiseret matematikhistorisk analyse af ikke lineær programmering: Udviklingshistorie og multipel opdagelse  
Ph.d.-afhandling af Tinne Hoff Kjeldsen
- 373/99 Criss-Cross Reduction of the Maslov Index and a Proof of the Yosida-Nicolaescu Theorem  
by: Bernhelm Booss-Bavnbek, Kenro Furutani and Nobukazu Otsuki
- 374/99 Det hydrauliske spring  
Et eksperimentelt studie af polygoner  
Og hastighedsprofiler  
Specialeafhandling af Anders Marcussen  
Vejledere: Tomas Bohr, Clive Ellegaard og Bent C. Jørgensen

- 375/99 Begrundelser for Matematikundervisningen  
i den lærde skole hhv. gymnasiet 1884-1914  
Historie-speciale af: Henrik Andreassen
- 376/99 Universality of AC conduction in  
disordered solids  
by: Jeppe C. Dyre, Thomas B. Schrøder
- 377/99 The Kuhn-Tucker Theorem in  
Nonlinear Programming:  
A Multiple Discovery?  
by: Tinne Hoff Kjeldsen
- .....
- 378/00 Solar energy preprints:  
Renewable energy sources and thermal energy storage  
Integration of photovoltaic cells into the global  
Energy system  
by: Bent Sørensen
- 379/00 EULERS DIFFERENTIALREGNING  
Eulers indførelse af differentialregningen stillet  
over for den moderne  
En tredjeseesters projektrapport på den  
naturvidenskabelige basisuddannelse  
af: Uffe Thomas Volmer Jankvist, Rie Rose Møller  
Pedersen, Maja Bagge Petersen  
Vejleder: Jørgen Larsen
- 380/00 Matematisk Modellering af Hjerterfunktionen  
Isovolumetrisk ventrikulær kontraktion og  
udpumpning til det kardiovaskulære system  
Speciale/3.moduls-rapport  
af: Gitte Andersen, Jakob Hilmer, Stine Weisbjerg  
Vejleder: Johnny Ottesen
- 381/00 Matematikviden og teknologiske kompetencer hos  
kortuddannede voksne  
- Rekognosceringer og konstruktioner i  
grænselandet mellem matematikkens  
didaktik og forskning i voksendannelse  
Ph.d.-afhandling af Tine Wedege
- 382/00 Den selvundvigende vandring  
Et matematisk professionsprojekt  
af: Martin Niss, Arnold Skimminge  
Vejledere: John Villumsen, Viggo Andreassen
- 383/00 Beviser i matematik  
af: Anne K.S.Jensen, Gitte M.Jensen,  
Jesper Thrane, Karen L.A.W. Wille,  
Peter Wulff  
Vejleder: Mogens Niss
- 384/00 Hopping in Disordered Media:  
A Model Glass Former and A Hopping Model  
Ph.d. thesis by Thomas B. Schrøder  
Supervisor: Jeppe C. Dyre
- 385/00 The Geometry of Cauchy Data Spaces  
by: Bernhelm Booss-Bavnbek, K. Furutani,  
K.P. Wojciechowski