

TEKST NR122

1986

ON THE MECHANISM OF GLASS IONIC CONDUCTIVITY

JEPPE C. DYRE
INSTITUTE OF MATHEMATICS AND PHYSICS
IMFUFA
UNIVERSITY OF ROSKILDE
POSTBOX 260
DK-4000 ROSKILDE
DENMARK

February 1986

TEKSTER fra

IMFUFA

ROSKILDE UNIVERSITETSCENTER
INSTITUT FOR STUDIET AF MATEMATIK OG FYSIK SAMT DERES
FUNKTIONER I UNDERVISNING, FORSKNING OG ANVENDELSER

Abstract

It is shown that most models can not explain the broad dielectric loss peaks and the correlation between dielectric loss and d.c. conductivity, which are both universal features of glass ionic conductivity. However, these features are reproduced by Stevels' and Taylor's old "random potential energy model" as is shown from a recent approximate solution of the model. It is argued that experimental data should preferably be presented in terms of the frequency-dependent conductivity instead of as dielectric loss peaks.

(Manuscript submitted to J. Non-Cryst. Solids)

Introduction

Despite several years of research the mechanism of glass ionic conductivity is still not understood (some reviews of the subject are found in refs. 1-7). In ionic conductive glasses one always observes both d.c. conductivity and dielectric relaxation. It is very interesting that these two phenomena are correlated, an observation of seemingly universal validity. If $\sigma(0)$ denotes the d.c. conductivity, Barton, Nakajima and Namikawa (BNN) discovered that most glasses obey [8,9,10]

$$\sigma(0) = p \epsilon_0 \Delta \epsilon \omega_m \quad (1)$$

where ϵ_0 is the vacuum permittivity, $\Delta \epsilon$ is the relaxation strength, ω_m is the loss peak frequency, and p is a temperature-independent numerical constant of order one. We remind that in media with a non-zero d.c. conductivity, the complex dielectric constant $\epsilon^* = \epsilon' - i \epsilon''$ is defined by [4]

$$\epsilon_0 \epsilon^*(\omega) = \frac{\sigma(\omega) - \sigma(0)}{i \omega} \quad (2)$$

where ω is the angular frequency and $\sigma = \sigma' + i \sigma''$ is the frequency-dependent conductivity. Equation (1) will be referred to as the BNN-relation. In particular it implies that $\sigma(0)$ and ω_m have the same activation energy. This was actually discovered before the BNN-relation, demonstrating that the dielectric relaxation is due to ionic motion [11]. In fig. 1 the

correlation between $\sigma(0)$ and $\epsilon_0 \Delta \epsilon \omega_m$ is examined for several glasses. It is clear that most glasses have a p-value of order one. The BNN-relation is thus experimentally very well-founded.

There is little doubt that the BNN-relation provides a key, perhaps the key, to a deeper understanding of ionic conductivity in glasses. Another observation of general validity is the very broad dielectric loss peaks. In this paper it will be shown that most current models are not able to explain both the BNN-relation and the broad loss peaks. In contrast, one of the oldest and simplest models works. This is Stevels' and Taylor's "random potential energy model" [2,12]. In this model, the loss peak frequency is simply the minimum ion jump frequency and the proportionality between $\sigma(0)$ and ω_m becomes physically obvious.

Critique of current models

Tomozawa has reviewed the most important models for the BNN-relation [1]. Three models predict simple Debye relaxation, namely Isard's, Charles' and Nakajima's model. In Isard's model the dielectric relaxation is assumed to be due to glass inhomogeneities which may derive e.g. from phase separation. If the different phases have different d.c. conductivity, dielectric relaxation of the bulk glass will result [13]. In Charles' model of alkali ionic conductivity in oxide glasses, the dielectric relaxation is due to defect jumps where a defect

is defined as a non-bridging oxygen atom surrounded by two alkali ions [14]. Finally, Nakajima has proposed a model for the BNN-relation which assumes that the ions are forced to follow certain random zig-zag paths frozen in the glass [9].

Because each of the above mentioned models predicts Debye relaxation, it is necessary to introduce a distribution of relaxation times into the models if the observed broad dielectric loss peaks are to be reproduced. Unfortunately, this is not possible without violating the BNN-relation. The reason is the following. Dielectric relaxation implies that the real part of the conductivity, $\sigma'(\omega)$, is an increasing function of frequency. For Debye relaxation with relaxation time τ , the increase in conductivity from low to high frequencies, $\Delta\sigma$, is easily found from eq. (2):

$$\Delta\sigma = \epsilon_0 \Delta\epsilon \tau^{-1} \quad (3)$$

The dielectric loss peak frequency is equal to τ^{-1} so from eqs. (1) and (3) one finds

$$\Delta\sigma = \frac{1}{\rho} \sigma(0) \quad (4)$$

Since $\sigma'(\omega)$ is monotonically increasing we therefore have

$$\sigma'(\omega) < (1 + \rho^{-1}) \sigma(0) \quad (5)$$

which is valid for all frequencies. Now, if the dielectric relaxation is assumed to be a sum of elementary Debye processes

each of which satisfies the BNN-relation, the resulting conductivity will still obey eq. (5) with a p of order one. But this is inconsistent with experiment because the broad loss peaks imply that $\sigma'(\omega) \gg \sigma(0)$ whenever $\omega \gg \omega_m$. The only way to save the models is to assume that the d.c. conductivities do not simply add algebraically. But then a whole new theory is needed for calculating $\sigma(0)$ and one may question the relevance of starting out with Debye processes each of which satisfies the BNN-relation. In conclusion, this procedure does not seem to work and one must look for a model which from the very beginning incorporates a broad distribution of relaxation times.

Tomozawa has proposed a model [1] based on Debye's and Falkenhagen's old theory of ionic solutions. This model is interesting because it has a broader loss peak than simple Debye relaxation. Still, the model predicts an excess conductivity ^{$\Delta\sigma$} of the same order of magnitude as $\sigma(0)$. This means that the loss peak is far from broad enough; if it is attempted to add several "Tomozawa" relaxation processes with different relaxation times one runs into the same problems as above, namely that the BNN-relation is violated.

A model which predicts a very broad dielectric loss peak is Doremus' model which is based on the space charge mechanism [5]. This model has $\epsilon''(\omega) \propto \omega^{-1/2}$ as $\omega \rightarrow \infty$ and a loss peak frequency which is related to the d.c. conductivity. However, because of the very nature of the model, the magnitude of the dielectric loss depends on sample dimensions while the BNN-relation is a true bulk property. Therefore, Doremus' model can

not explain the BNN-relation [1].

Stevens' and Taylor's "random potential energy model" [2,12] is perhaps the oldest model. In the next section it will be shown that this model is able to explain both the BNN-relation and the broad loss peaks.

The random potential energy model

In this model it is assumed that the ions feel a more or less random potential energy deriving from the random network structure of the glass. An ion spends most time at potential energy minima, but occasionally it gains energy by thermal fluctuations to pass the energy barrier which separates adjacent potential energy minima. The ion jump frequency, γ , depends on the energy barrier, ΔE , as

$$\gamma = \gamma_0 e^{-\Delta E/kT} \quad (6)$$

where γ_0 is the attempt frequency (usually $\approx 10^{12}$ Hz), k is the Boltzmann constant, and T is the temperature. Because of the randomness of the potential energy surface all energy barriers are equally likely, so from eq. (6) it is concluded that the jump frequency probability distribution, $p(\gamma)$, varies as

$$p(\gamma) \propto \gamma^{-1}. \quad (7)$$

If the ions can jump to more than one nearest neighbour energy

minimum, the resulting $p(\gamma)$ changes only insignificantly from eq. (7) which forms the basis of the present treatment of Stevels' and Taylor's model.

The model of a random medium with the jump frequency distribution of eq. (7) has recently been considered by the author [15,16]. The treatment makes use of the CTRW approximation [17]. In order to normalize $p(\gamma)$ it is necessary to introduce two cut-off's, γ_{min} and γ_{max} . The existence of γ_{min} in real glasses is obvious: if $\gamma_{min} = 0$ a zero d.c. conductivity results. On the other hand, experimentally there is no sign of any maximum jump frequency so γ_{max} should be put equal to γ_0 . Since laboratory frequencies are always much smaller than γ_0 , it is convenient to let γ_0 go to infinity. In this limit one finds [15]

$$\sigma(\omega) = \sigma(0) \frac{i\omega\tau}{\ln(1+i\omega\tau)} \quad (8)$$

where

$$\tau = \gamma_{min}^{-1} \quad (9)$$

Equation (8) does not, of course, include the contribution to the conductivity from the high frequency dielectric constant. Substituting now eq. (8) into eq. (2) we get

$$\epsilon_0 \Delta \epsilon = \frac{1}{2} \sigma(0) \tau \quad (10)$$

and for the dielectric loss

$$\epsilon''(\omega) = 2 \Delta \epsilon \left[\frac{\arctan(\omega\tau)}{(\ln\sqrt{1+(\omega\tau)^2})^2 + (\arctan(\omega\tau))^2} - \frac{1}{\omega\tau} \right] \quad (11)$$

From eq. (11) the loss peak frequency can easily be determined numerically. It is given by $\omega_m \tau = 4,71$ which corresponds to a p-value of the BNN-relation of

$$p = 0,42. \quad (12)$$

Since 0,42 is a numerical constant of order one, the BNN-relation is satisfied by the random potential energy model.

The model predicts a very broad dielectric loss peak as is clear from fig. 2. In this figure the model prediction is compared to experimental data from a typical sodium silicate glass. There is a qualitative agreement between theory and experiment.

Discussion

The derivation of eq. (8) makes use of a number of simplifying assumptions: 1) All ion-ion interactions are ignored, 2) all energy barriers are assumed to be equally likely in the interval from zero to a sharply defined maximum energy barrier, and 3) the model is solved within the CTRW

approximation which is the simplest possible non-trivial mean-field approximation for calculating $\sigma(\omega)$ in random media. The validity of these approximations will not be critically examined here. One point is perhaps worth noticing. The present use of the CTRW approximation (letting $\gamma_0 \rightarrow \infty$) is formally equivalent to the electric modulus approach of Macedo, Moynihan and Bose [18]. The physical interpretation of this analogy is not quite clear at present [16,19]. One possibility is that the electrical equivalent circuit of ref. 18 (fig.3) reflects a one-dimensional character for conduction processes in glasses [19]. Another possibility is to regard the electric modulus approach as the relevant description when inhomogeneities are present, i.e. as a generalization of Isard's model [16,18]. Because of the analogy to the CTRW, if the inhomogeneities have d.c. conductivities with a probability distribution $p(\sigma(0)) \propto \sigma(0)^{-1}$, the bulk frequency-dependent conductivity is again given by eq. (8). As regards electrical properties, the random potential energy model is therefore indistinguishable from a generalized Isard model (if this model really is faithfully represented by the equivalent circuit of fig. 3).

For a number of years it has been thought that the random potential energy model, while appealing, can not possibly be correct [1,6]. The BNN-relation implies that the activation energies of $\sigma(0)$ and ω_m are equal, and this has been interpreted as implying that the a.c. conductivity must be due to ion jumps with the same activation energy as the d.c. conductivity. The argument is incorrect, however. As is clear from the preceding section, while the activation energy of

$\sigma(\omega)$ is certainly equal to that of χ_{min} , there is absolutely no problem with having larger jump frequencies involved in the conduction process, corresponding to smaller energy barriers. Another common objection to Stevels' and Taylor's model is that no model based on a distribution of energy barriers can predict a loss peak with a temperature-independent shape. This is wrong; in the present model the jump frequency distribution is given by eq. (7) for all temperatures and therefore the shape of the loss peak indeed is temperature-independent.

Glass ionic conductivity data are usually presented in terms of the dielectric loss when the frequency dependence is discussed. However, because d.c. conductivity and dielectric loss are both due to ionic motion, the subtraction of $\sigma(\omega)$ in eq. (2) really has no physical justification. In the author's opinion, data should preferably be presented in terms of the real part of the conductivity. As an example, in fig. 4 the data of fig. 2 are shown together with model predictions (eq. 8). A number of important features of glass ionic conductivity become immediately clear when data are presented this way. The characteristic frequency which marks the onset of a.c. conductivity is just the dielectric loss peak frequency ω_m . On a time-scale larger than ω_m^{-1} , $\sigma(\omega)$ is frequency-independent so the glass "looks" homogeneous to the ions. On a time-scale smaller than ω_m^{-1} , $\sigma(\omega)$ is strongly frequency-dependent. The obvious physical interpretation is that ω_m is the minimum ion jump frequency; then many jumps must be involved for times $\gg \omega_m^{-1}$. In order to account for the strong

increase of $\sigma'(\omega)$ for $\omega > \omega_m$, jump frequencies much larger than ω_m must however also be present in the glass. The random potential energy model has exactly these features. In this model a wide range of jump frequencies is present, deriving from a range of activation energies.

The spread in activation energies in Stevels' and Taylor's model is not an arbitrary postulate. On the contrary, it is a simple fact which follows immediately when experiments are presented in terms of the frequency-dependent conductivity in a log-log plot as in fig. 4 but without the normalization: The shape of the conductivity curve is always temperature-independent. Because of the BNN-relation, when the temperature changes the conductivity curve is simply displaced in the direction 45° to the positive x- and y-axis. It is now clear by inspection that the a.c. conductivity indeed has a smaller activation energy than the d.c. conductivity. And from the curvature of the conductivity curve it can be concluded that a whole range of activation energies must be involved. Thus, the experimental facts themselves apparently forces one to adopt Stevels' and Taylor's approach and base the theory of glass ionic conductivity on a distribution of energy barriers.

Another reason for preferring to discuss glass ionic conductivity in terms of $\sigma(\omega)$, is the conceptual dilemma involved in the use of $\xi(\omega)$ as defined by eq. (2). One would like to interpret $\xi(\omega)$ in terms of a dielectric polarisation but it is difficult, if not impossible, to tell which ion jumps contribute to $\sigma(0)$ and which contribute to $\xi(\omega)$. This dilemma is old and well-known. It derives from the fact that it

is not possible to distinguish between bound and free charges in a.c. electric fields [20]. The problem disappears when one operates only with $\sigma(\omega)$.

Conclusions

Most current models of glass ionic conductivity can not explain both the BNN-relation and the observed broad dielectric loss peaks. On the other hand, the random potential energy model is rather successful. This model involves no ad hoc assumptions and it is probably the simplest possible model which is not totally unrealistic. The model is based on a fact which can be deduced directly from experiment when presented in terms of the frequency-dependent conductivity, namely the existence of a distribution of energy barriers. The BNN-relation is satisfied by the model. Here, the relation is nothing but the statement that the d.c. conductivity is proportional to the minimum ion jump frequency. The jump frequency distribution eq. (7) strongly emphasizes the smallest jump frequencies which are also the most important for $\sigma(0)$ because they partially act as traps, so this proportionality is by no means surprising. In addition to the BNN-relation, the model also reproduces the observed broad asymmetric loss peaks, the temperature independence of the loss peak shape, and the almost frequency-independent dielectric loss as $\omega \rightarrow \infty$. These are all well-known characteristics of glass ionic conductivity [1,4]. The quantitative model predictions are not entirely successful, though (fig. 2 and fig.

4). This is undoubtedly because it is too simple to regard all energy barriers as equally likely. Refinements of the model must make use of a more realistic energy barrier distribution.

Admittedly, this introduces a temperature dependence into the loss peak shape, but in many cases the effect is only weak and may still be consistent with experiment. The real challenge for the future is to make a coherent theory in which the energy barrier distribution is correlated to the glass structure and conditions of glass formation.

References

1. M. Tomozawa, in: "Treatise on Materials Science", Vol. 12, ed. M. Tomozawa (Academic Press, New York, 1977) p.283.
2. J. M. Stevels, in: "Handbuch der Physik", Vol. 20, ed. S. Flügge (Springer Verlag, Berlin, 1957) p. 350.
3. K. Hughes and J. O. Isard, in: "Physics of Electrolytes", Vol. 1, ed. J. H. Hladik (Academic Press, New York, 1972) p. 351.
4. A. E. Owen, J. Non-Cryst. Solids 25 (1977) 372.
5. R. H. Doremus, "Glass Science" (Wiley, New York, 1973).
6. D. Ravaine and J. L. Souquet, in: "Solid Electrolytes", ed. P. Hagenmuller and W. Van Gool (Academic Press, New York, 1978) p. 277.
7. K. J. Rao and R. Parthasarathy, Proc. Indian Acad. Sci. (Chem. Sci.) 94 (1985) 201.
8. J. L. Barton, Verres et Refr. 20 (1966) 328.
9. T. Nakajima, in: "1971 Annual Report, Conference on Electric Insulation and Dielectric Phenomena", Nat. Acad. of Sci.
10. H. Namikawa, J. Non-Cryst. Solids 18 (1975) 173.
11. H. E. Taylor, Trans. Farad. Soc. 52 (1956) 873.
12. H. E. Taylor, J. Soc. Glass Technol. 41 (1957) 350T.
13. J. O. Isard, Proc. Inst. Elec. Eng. Suppl. 22, 109 Part B (1962) 440.
14. R. J. Charles, J. Appl. Phys. 32 (1961) 1115.
15. J. C. Dyre, Phys. Lett. 108A (1985) 457.
16. J. C. Dyre, J. Physique Colloq. 46, C-8 (1985) 343.

17. H. Scher and M. Lax, Phys. Rev. B7 (1973) 4491.
18. P. B. Macedo, C. T. Moynihan and R. Bose, Phys. Chem. Glasses 13 (1972) 171.
19. J. C. Dyre, IMFUFA tekst no. 97 (Univ. of Roskilde, 1985).
20. N. W. Ashcroft and N. D. Mermin, "Solid State Physics" (Holt, Rinehart & Winston, New York, 1976) appendix K.

Figure Captions

- Fig. 1 : Test of the BNN-relation for several glasses (note that $2\pi f_m = \omega_m$) (reproduced from ref. 9).
- Fig. 2 : Model prediction for the dielectric loss (full curve) and experimental data from a typical sodium silicate glass (data taken from ref. 4). The dashed curve is the Debye dielectric loss peak.
- Fig. 3 : Equivalent circuit of the CTRW approximation in the $\chi_0 \rightarrow \infty$ limit [16]. The circuit was suggested by Macedo, Moynihan and Bose in their electric modulus approach towards glass ionic conductivity [18].
- Fig. 4 : Model prediction for the real part of the conductivity (full curve) and the experimental data of fig. 2. For the data points the conductivity has been calculated assuming $p=1$ in the BNN-relation and utilizing the Kramers-Kronig relation for calculating $\Delta \epsilon$ ($= 3,7 \cdot \epsilon''_{\max}$). The characteristic time τ is a fitting parameter for the data points in this figure.




Fig. 1




fig.2




fig. 3




fig. 4

- 1/78 "TANKER OM EN PRAKSIS" - et matematikprojekt.
 Projektrapport af Anne Jensen, Lena Lindenskov, Marianne Kesselhahn og Nicolai Lomholt.
 Vejleder: Anders Madsen.
- 2/78 "OPTIMERING" - Menneskets forøgede beherskelsesmuligheder af natur og samfund.
 Projektrapport af Tom J. Andersen, Tommy R. Andersen, Gert Kreinøe og Peter H. Lassen.
 Vejleder: Bernhelm Booss.
- 3/78 "OPGAVESAMLING", breddekursus i fysik. Nr. 3 er a jour ført i marts 1984
 Lasse Rasmussen, Aage Bonde Kræmmer, Jens Højgaard Jensen.
- 4/78 "TRE ESSAYS" - om matematikundervisning, matematiklæreruddannelsen og videnskabsrindalismen. Nr. 4 er p.t. udgået.
 Mogens Niss.
- 5/78 "BIBLIOGRAFISK VEJLEDNING til studiet af DEN MODERNE FYSIKS HISTORIE". Nr. 5 er p.t. udgået.
 Helge Kragh.
- 6/78 "NOGLE ARTIKLER OG DEBATINDLÆG OM - læreruddannelse og undervisning i fysik, og - de naturvidenskabelige fags situation efter studenteroprøret".
 Karin Beyer, Jens Højgaard Jensen og Bent C. Jørgensen.
- 7/78 "MATEMATIKKENS FORHOLD TIL SAMFUNDSØKONOMIEN". Nr. 7 er udgået.
 B.V. Gnedenko.
- 8/78 "DYNAMIK OG DIAGRAMMER". Introduktion til energy-bond-graph formalismen.
 Peder Voetmann Christiansen.
- 9/78 "OM PRAKSIS' INDFLYDELSE PÅ MATEMATIKKENS UDVIKLING". - Motiver til Kepler's: "Nova Stereometria Doliorum Vinarioum".
 Projektrapport af Lasse Rasmussen.
 Vejleder: Anders Madsen.
-
- 10/79 "TERMODYNAMIK I GYMNASIET".
 Projektrapport af Jan Christensen og Jeanne Mortensen.
 Vejledere: Karin Beyer og Peder Voetmann Christiansen.
- 11/79 "STATISTISKE MATERIALER"
 red. Jørgen Larsen
- 12/79 "LINEÆRE DIFFERENTIALLIGNINGER OG DIFFERENTIALLIGNINGSSYSTEMER". Nr. 12 er udgået
 Mogens Brun-Heefelt
- 13/79 "CAVENDISH'S FORSØG I GYMNASIET".
 Projektrapport af Gert Kreinøe.
 Vejleder: Albert Chr. Paulsen

- 14/79 "BOOKS ABOUT MATHEMATICS: History, Philosophy, Education, Models, System Theory, and Works of Reference etc. A Bibliography".
Else Høyrup.
Nr. 14 er p.t. udgået.
- 15/79 "STRUKTUREL STABILITET OG KATASTROFER i systemer i og udenfor termodynamisk ligevægt".
Specialeopgave af Leif S. Striegler.
Vejleder: Peder Voetmann Christiansen.
- 16/79 "STATISTIK I KRÆFTFORSKNINGEN".
Projektrapport af Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen.
- 17/79 "AT SPØRGE OG AT SVARE i fysikundervisningen".
Albert Christian Paulsen.
- 18/79 "MATHEMATICS AND THE REAL WORLD", Proceedings of an International Workshop, Roskilde University Centre, Denmark, 1978. Preprint.
Bernhelm Booss & Mogens Niss (eds.).
- 19/79 "GEOMETRI, SKOLE OG VIRKELIGHED".
Projektrapport af Tom J. Andersen, Tommy R. Andersen og Per H.H. Larsen.
Vejleder: Mogens Niss.
- 20/79 "STATISTISKE MODELLER TIL BESTEMMELSE AF SIKRE DOSER FOR CARCINOGENE STOFFER".
Projektrapport af Michael Olsen og Jørn Jensen.
Vejleder: Jørgen Larsen.
- 21/79 "KONTROL I GYMNASIET - FORMAL OG KONSEKVENSER".
Projektrapport af Crilles Bacher, Per S. Jensen, Preben Jensen og Torben Nysteen.
- 22/79 "SEMIOTIK OG SYSTEMEGENSKABER (1)".
1-port lineært response og støj i fysikken.
Peder Voetmann Christiansen.
- 23/79 "ON THE HISTORY OF EARLY WAVE MECHANICS - with special emphasis on the role of reality".
-
- 24/80 "MATEMATIKOPFATTELSE HOS 2.G'ERE".
a+b 1. En analyse. 2. Interviewmateriale.
Projektrapport af Jan Christensen og Knud Lindhardt Rasmussen.
Vejleder: Mogens Niss.
Nr. 24 a+b er p.t. udgået.
- 25/80 "EKSAMENSOPGAVER", Dybdemodul/fysik 1974-79.
- 26/80 "OM MATEMATISKE MODELLER".
En projektrapport og to artikler.
Jens Højgaard Jensen m.fl.
- 27/80 "METHODOLOGY AND PHILOSOPHY OF SCIENCE IN PAUL DIRAC'S PHYSICS".
Helge Kragh.
- 28/80 "DIELEKTRISK RELAXATION - et forslag til en ny model bygget på væskernes viscoelastiske egenskaber".
Projektrapport, speciale i fysik, af Gert Kreinøe.
Vejleder: Niels Boye Olsen.

- 29/80 "ODIN - undervisningsmateriale til et kursus i differentialligningsmodeller".
 Projekt rapport af Tommy R. Andersen, Per H.H. Larsen og Peter H. Lassen.
 Vejleder: Mogens Brun Heefelt
- 30/80 "FUSIONSENERGIEN - - - ATOMSAMFUNDETS ENDESTATION".
 Oluf Danielsen. Nr. 30 er udgået.
 Udkommer medio 1982 på Fysik-, Matematik- og Kemilærernes forlag.
- 31/80 "VIDENSKABSTEORETISKE PROBLEMER VED UNDERVISNINGSSYSTEMER BASERET PÅ MÆNGDELÆRE".
 Projekt rapport af Troels Lange og Jørgen Karrebæk.
 Vejleder: Stig Andur Pedersen. Nr. 31 er p.t. udgået
- 32/80 "POLYMERE STOFFERS VISCOELASTISKE EGENSKABER - BELYST VED HJÆLP AF MEKANISKE IMPEDANSMÅLINGER OG MOSSBAUER-EFFEKTMÅLINGER".
 Projekt rapport, speciale i fysik, af Crilles Bacher og Preben Jensen.
 Vejledere: Niels Boye Olsen og Peder Voetmann Christiansen.
- 33/80 "KONSTITUERING AF FAG INDEN FOR TEKNISK-NATURVIDENSKABELIGE UDDANNELSER. I-II".
 Arne Jakobsen.
- 34/80 "ENVIRONMENTAL IMPACT OF WIND ENERGY UTILIZATION".
 ENERGY SERIES NO.1. Nr. 34 er udgået.
 Bent Sørensen. Publ. i "Renewable Sources of Energy and the Environment", Tycooli International Press, Dublin, 1981.
- 35/80 "HISTORISKE STUDIER I DEN NYERE ATOMFYSIKS UDVIKLING".
 Helge Kragh.
- 36/80 "HVAD ER MENINGEN MED MATEMATIKUNDERVISNINGEN ?".
 Fire artikler.
 Mogens Niss.
- 37/80 "RENEWABLE ENERGY AND ENERGY STORAGE".
 ENERGY SERIES NO.2.
 Bent Sørensen.
-
- 38/81 "TIL EN HISTORIETEORI OM NATURERKENDELSE, TEKNOLOGI OG SAMFUND".
 Projekt rapport af Erik Gade, Hans Hedal, Henrik Lau og Finn Physant. Nr. 38 er p.t. udgået
 Vejledere: Stig Andur Pedersen, Helge Kragh og Ib Thiersen.
- 39/81 "TIL KRITIKKEN AF VÆKSTØKONOMIEN".
 Jens Højgaard Jensen.
- 40/81 "TELEKOMMUNIKATION I DANMARK - oplæg til en teknologivurdering".
 Projekt rapport af Arne Jørgensen, Bruno Petersen og Jan Vedde. Nr. 40 er p.t. udgået
 Vejleder: Per Nørgaard.
- 41/81 "PLANNING AND POLICY CONSIDERATIONS RELATED TO THE INTRODUCTION OF RENEWABLE ENERGY SOURCES INTO ENERGY SUPPLY SYSTEMS".
 ENERGY SERIES NO.3.
 Bent Sørensen.

- 42/81 "VIDENSKAB TEORI SAMFUND - En introduktion til materialistiske videnskabsopfattelser".
Helge Kragh og Stig Andur Pedersen.
- 43/81 1. "COMPARATIVE RISK ASSESSMENT OF TOTAL ENERGY SYSTEMS".
2. "ADVANTAGES AND DISADVANTAGES OF DECENTRALIZATION".
ENERGY SERIES NO.4.
Bent Sørensen.
- 44/81 "HISTORISK UNDERSØGELSE AF DE EKSPERIMENTELLE FORUDSÆTNINGER FOR RUTHERFORDS ATOMMODEL".
Projektrapport af Niels Thor Nielsen.
Vejleder: Bent C. Jørgensen.
-

45/82

- 46/82 "EKSEMPLARISK UNDERVISNING OG FYSISK ERKENDELSE - I+II ILLUSTRERET VED TO EKSEMPLER".
Projektrapport af Torben O. Olsen, Lasse Rasmussen og Niels Dreyer Sørensen.
Vejleder: Bent C. Jørgensen.
- 47/82 "BARSEBÄCK OG DET VÆRST OFFICIELT-TÆNKELIGE UHELD".
ENERGY SERIES NO.5.
Bent Sørensen.
- 48/82 "EN UNDERSØGELSE AF MATEMATIKUNDERVISNINGEN PÅ ADGANGSKURSUS TIL KØBENHAVNS TEKNIKUM".
Projektrapport af Lis Eilertzen, Jørgen Karrebæk, Troels Lange, Preben Nørregaard, Lissi Pedersen, Laust Rishøj, Lill Røn, Isac Showiki.
Vejleder: Mogens Niss.
- 49/82 "ANALYSE AF MULTISPEKTRALE SATELLITBILLEDER".
Projektrapport af Preben Nørregaard.
Vejledere: Jørgen Larsen & Rasmus Ole Rasmussen.
- 50/82 "HERSLEV - MULIGHEDER FOR VEDVARENDE ENERGI I EN LANDSBY". ENERGY SERIES NO.6.
Rapport af Bent Christensen, Bent Hove Jensen, Dennis B. Møller, Bjarne Laursen, Bjarne Lillethorup og Jacob Mørch Pedersen.
Vejleder: Bent Sørensen.
- 51/82 "HVAD KAN DER GØRES FOR AT AFHJÆLPE PIGERS BLOKERING OVERFOR MATEMATIK?"
Projektrapport af Lis Eilertzen, Lissi Pedersen, Lill Røn og Susanne Stender.
- 52/82 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS"
Bernhelm Booss & Krzysztof Wojciechowski.
- 53/82 "THE CONSTITUTION OF SUBJECTS IN ENGINEERING EDUCATION".
Arne Jakobsen & Stig Andur Pedersen.
- 54/82 "FUTURES RESEARCH" - A Philosophical Analysis of Its Subject-Matter and Methods.
Stig Andur Pedersen & Johannes Witt-Hansen.

- 55/82 "MATEMATISKE MODELLER" - Litteratur på Roskilde
Universitetsbibliotek. Vedr. tekst nr. 55/82:
Se også tekst 62/83.
En bibliografi.
Else Høyrup.
- 56/82 "ÉN - TO - MANGE" -
En undersøgelse af matematisk økologi.
Projektrapport af Troels Lange.
Vejleder: Anders Madsen.
-
- 57/83 "ASPECT EKSPERIMENTET" - Nr. 57 er udgået.
Skjulte variable i kvantemekanikken?
Projektrapport af Tom Juul Andersen.
Vejleder: Peder Voetmann Christiansen.
- 58/83 "MATEMATISKE VANDRINGER" - Modelbetragtninger
over spredning af dyr mellem småbiotoper i
agerlandet.
Projektrapport af Per Hammershøj Jensen &
Lene Vagn Rasmussen.
Vejleder: Jørgen Larsen.
- 59/83 "THE METHODOLOGY OF ENERGY PLANNING".
ENERGY SERIES NO. 7.
Bent Sørensen.
- 60/83 "MATEMATISK MODEKSPERTISE" - et eksempel.
Projektrapport af Erik O. Gade, Jørgen Karrebæk og
Preben Nørregaard.
Vejleder: Anders Madsen.
- 61/83 "FYSIKS IDEOLOGISKE FUNKTION", som et eksempel på
en naturvidenskab - historisk set.
Projektrapport af Annette Post Nielsen.
Vejledere: Jens Høyrup, Jens Højgaard Jensen og
Jørgen Vogelius.
- 62/83 "MATEMATISKE MODELLER" - Litteratur på Roskilde
Universitetsbibliotek.
En bibliografi. 2. rev. udgave
Else Høyrup
- 63/83 "CREATING ENERGY FUTURES: A SHORT GUIDE TO
ENERGY PLANNING".
ENERGY SERIES No. 8
David Crossley & Bent Sørensen
- 64/83 "VON MATHEMATIK UND KRIEG".
Bernhelm Booss og Jens Høyrup.
- 65/83 "ANVENDT MATEMATIK - TEORI ELLER PRAKSIS".
Projektrapport af Per Hedegård Andersen, Kirsten
Habekost, Carsten Holst-Jensen, Annelise von Moos,
Else Marie Pedersen, Erling Møller Pedersen.
Vejledere: Bernhelm Booss & Klaus Grünbaum
- 66/83 "MATEMATISKE MODELLER FOR PERIODISK SELEKTION I
ESCHERICHIA COLI".
Projektrapport af Hanne Lisbet Andersen, Ole
Richard Jensen og Klavs Frisdahl.
Vejledere: Jørgen Larsen og Anders Hede Madsen

- 67/83 "ELIPSOIDE METODEN - EN NY METODE TIL LINEÆR PROGRAMMERING?"
Projektrapport af Lone Biilmann og Lars Boye
Vejleder: Mogens Brun Heefelt
- 68/83 "STOKASTISKE MODELLER I POPULATIONSGENETIK"
- til kritikken af teoriladede modeller.
Projektrapport af Lise Odgård Gade, Susanne Hansen, Michael Hviid, Frank Mølgård Olsen.
Vejleder: Jørgen Larsen.
- 69/83 "ELEVFORUDSÆTNINGER I FYSIK"
- en test i l.g med kommentarer
Albert Chr. Paulsen
- 70/83 "INDLÆRINGS- OG FORMIDLINGSPROBLEMER I MATEMATIK PÅ VOKSEUNDERVISNINGSNIVEAU"
Projektrapport af Hanne Lisbet Andersen, Torben J. Andreasen, Svend Åge Houmann, Helle Glerup Jensen, Keld Fl. Nielsen, Lene Vagn Rasmussen.
Vejleder: Klaus Grünbaum & Anders H. Madsen
- 71/83 "PIGER OG FYSIK"
- et problem og en udfordring for skolen?
Karin Beyer, Sussanne Blegaa, Birthe Olsen, Jette Reich & Mette Vedelsby
- 72/83 "VERDEN IFØLGE PEIRCE" - to metafysiske essays, om og af C.S. Peirce.
Peder Voetmann Christiansen
- 73/83 "EN ENERGIANALYSE AF LANDBRUG"
- økologisk contra traditionelt
ENERGY SERIES No. 9
Specialeopgave i fysik af Bent Hove Jensen
Vejleder: Bent Sørensen
-
- 74/84 "MINIATURISERING AF MIKROELEKTRONIK" - om videnskabeliggjort teknologi og nytten af at lære fysik
Projektrapport af Bodil Harder og Linda Szkotak Jensen.
Vejledere: Jens Højgaard Jensen og Bent C. Jørgensen
- 75/84 "MATEMATIKUNDERVISNINGEN I FREMTIDENS GYMNASIUM"
- Case: Lineær programmering
Projektrapport af Morten Blomhøj, Klavs Frisdahl, Frank Mølgaard Olsen
Vejledere: Mogens Brun Heefelt & Jens Bjørneboe
- 76/84 "KERNEKRAFT I DANMARK?" - Et høringssvar indkaldt af miljøministeriet, med kritik af miljøstyrelsens rapporter af 15. marts 1984.
ENERGY SERIES No. 10
Af Niels Boye Olsen og Bent Sørensen
- 77/84 "POLITISKE INDEKS - FUP ELLER FAKTA?"
Opinionsundersøgelser belyst ved statistiske modeller
Projektrapport af Svend Åge Houmann, Keld Nielsen, Susanne Stender
Vejledere: Jørgen Larsen & Jens Bjørneboe

- 78/84 "JÆVNSTRØMSLEDNINGSEVNE OG GITTERSTRUKTUR I AMORFT GERMANIUM"
Specialerapport af Hans Hedal, Frank C. Ludvigsen og Finn C. Physant
Vejleder: Niels Boye Olsen
- 79/84 "MATEMATIK OG ALMENDANNELSE"
Projektrapport af Henrik Coster, Mikael Wennerberg Johansen, Povl Kattler, Birgitte Lydholm og Morten Overgaard Nielsen.
Vejleder: Bernhelm Booss
- 80/84 "KURSUSMATERIALE TIL MATEMATIK B"
Mogens Brun Heefelt
- 81/84 "FREKVENSAFHÆNGIG LEDNINGSEVNE I AMORFT GERMANIUM"
Specialerapport af Jørgen Wind Petersen og Jan Christensen
Vejleder: Niels Boye Olsen
- 82/84 "MATEMATIK- OG FYSIKUNDERVISNINGEN I DET AUTOMATISEREDE SAMFUND"
Rapport fra et seminar afholdt i Hvidovre 25-27 april 1983
Red.: Jens Højgaard Jensen, Bent C. Jørgensen og Mogens Niss
- 83/84 "ON THE QUANTIFICATION OF SECURITY" nr. 83 er p.t. udgået
PEACE RESEARCH SERIES NO. 1
af Bent Sørensen
- 84/84 "NOGLE ARTIKLER OM MATEMATIK, FYSIK OG ALMENDANNELSE".
Jens Højgaard Jensen, Mogens Niss m. fl.
- 85/84 "CENTRIFUGALREGULATORER OG MATEMATIK"
Specialerapport af Per Hedegård Andersen, Carsten Holst-Jensen, Else Marie Pedersen og Erling Møller Pedersen
Vejleder: Stig Andur Pedersen
- 86/84 "SECURITY IMPLICATIONS OF ALTERNATIVE DEFENSE OPTIONS FOR WESTERN EUROPE"
PEACE RESEARCH SERIES NO. 2
af Bent Sørensen
- 87/84 "A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY IN DISORDERED SOLIDS"
af Jeppe C. Dyre
- 88/84 "RISE, FALL AND RESURRECTION OF INFINITESIMALS"
af Detlef Laugwitz
- 89/84 "FJERNVARMEOPTIMERING"
af Bjarne Lillethorup & Jacob Mørch Pedersen
- 90/84 "ENERGI I 1.G. en teori for tilrettelæggelse"
af Albert Chr. Paulsen
-
- 91/85 "KVANTETEORI FOR GYMNASIET"
1. Lærervejledning
Projektrapport af: Birger Lundgren, Henning Sten Hansen og John Johansson
Vejleder: Torsten Meyer

- 92/85 "KVANTETEORER FOR GYMNASIET
2. Materiale
Projektrapport af: Birger Lundgren, Henning
Sten Hansen og John Johansson
Vejleder: Torsten Meyer
- 93/85 "THE SEMIOTICS OF QUANTUM-NON-LOCALITY"
af Peder Voetmann Christiansen
- 94/85 "TREENIGHEDEN BOURBAKI - generalen, matematikeren
og ånden"
Projektrapport af: Morten Blomhøj, Klavs Frisdahl
og Frank M. Olsen
Vejleder: Mogens Niss
- 95/85 "AN ALTERNATIV DEFENSE PLAN FOR WESTERN EUROPE"
Peace research series no. 3
af Bent Sørensen
- 96/85 "ASPEKTER VED KRAFTVARMEFORSYNING"
af Bjarne Lillethorup
Vejleder: Bent Sørensen
- 97/85 "ON THE PHYSICS OF A.C. HOPPING CONDUCTIVITY"
Jeppe C. Dyre
- 98/85 "VALGMULIGHEDER I INFORMATIONSSALDEREN"
af Bent Sørensen
- 99/85 "Der er langt fra Q til R"
Projektrapport af: Niels Jørgensen og Mikael Klintorp
Vejleder: Andur Pedersen
- 100/85 "TALSYSTEMETS OPBYGNING"
af Mogens Niss
- 101/85 "EXTENDED MOMENTUM THEORY FOR WINDMILLS
IN PERTURBATIVE FORM"
af Ganesh Sengupta
- 102/85 "OPSTILLING OG ANALYSE AF MATEMATISKE MODELLER, BELYST VED
MODELLER OVER KØERS FODEROPTAGELSE OG - OMSÆTNING"
Projektrapport af: Lis Eilertzen, Kirsten Habekost, Lill Røn
og Susanne Stender
Vejleder: Klaus Grünbaum
- 103/85 "ØDSLE KOLDKRIGERE & VIDENSKABENS LYSE IDEER"
Projektrapport af: Niels Ole Dam og Kurt Jensen
Vejleder: Bent Sørensen
- 104/85 "ANALOGREGNEMASKINEN OG LORENZLIGNINGER"
af: Jens Jæger
- 105/85 "THE FREQUENCY DEPENDENCE OF THE SPECIFIC HEAT AT THE GLASS
TRANSITION"
af Tage Christensen
"A SIMPLE MODEL OF AC HOPPING CONDUCTIVITY"
af Jeppe C. Dyre
Contributions to the Third International Conference on the
Structure of Non-Crystalline Materials held in Grenoble
July 1985
- 106/85 "QUANTUM THEORY OF EXTENDED PARTICLES"
af Bent Sørensen
- 107/85 "ÉN MYG GØR INGEN EPIDEMI"
- flodblindhed som eksempel på matematisk modellering af et
epidemiologisk problem.
Projektrapport af: Per Hedegård Andersen, Lars Boye, Carsten

Holst Jensen, Else Marie Pedersen og Erling Møller Pedersen
Vejleder: Jesper Larsen

108/85 "APPLICATIONS AND MODELLING IN THE MATHEMATICS CURRICULUM"

- state and trends -
af Mogens Niss

109/85 "COX I STUDIETIDEN"

- Cox's regressionsmodel anvendt på studenteroplysninger
fra RUC
Projektrapport af: Mikael Wennerberg Johansen, Poul Kattler
og Torben J. Andreasen
Vejleder: Jørgen Larsen

110/85 " PLANNING FOR SECURITY "

af Bent Sørensen

111/85 "JORDEN RUNDT PÅ FLADE KORT"

Projektrapport af: Birgit Andresen, Beatriz Quinones og
Jimmy Staal

Vejleder: Mogens Niss

112/85 "VIDENSKABELIGGØRELSE AF DANSK TEKNOLOGISK INNOVATION
FREM TIL 1950 - BELYST VED EKSEMPLER"

Projektrapport af: Erik Odgaard Gade, Hans Hedal, Frank
C. Ludvigsen, Annette Post Nielsen og Finn Physant.
Vejleder: Claus Bryld og Bent C. Jørgensen

113/85 "DESUSPENSION OF SPLITTING ELLIPTIC SYMBOLS II"

Af: Bernhelm Booss og Krzysztof Wojciechowski

114/85 "ANVENDELSE AF GRAFISKE METODER TIL ANALYSE AF
KONTIGENSTABELLER"

Projektrapport af: Lone Billmann, Ole R. Jensen
og Anne-Lise von Moos
Vejleder: Jørgen Larsen

115/85 "MATEMATIKKENS UDVIKLING OP TIL RENESSANCEN"

af: Mogens Niss

116/85 "A PHENOMENOLOGICAL MODEL FOR THE MEYER-
NELDEL RULE"

af: Jeppe C. Dyre

117/85 "KRAFT & FJERNVARMEOPTIMERING"

af: Jacob Mørch Pedersen
Vejleder: Bent Sørensen

118/85 "TILFÆLDIGHEDEN OG NØDVENDIGHEDEN IFØLGE
PEIRCE OG FYSIKKEN"

af: Peder Voetmann Christiansen

119/86 "DET ER GANSKE VIST-- EUKLIDS FEMTE POSTULAT
KUNNE NOK SKABE RØRE I ANDEDAMMEN"

af: Iben Maj Christiansen
Vejleder: Mogens Niss

120/86 "ET ANTAL STATISTISKE STANDARDMODELLER"

af: Jørgen Larsen